

AD-A113 192

CALIFORNIA UNIV LOS ANGELES DEPT OF CHEMISTRY

F/G 11/9

REACTIONS AT THE METAL VERTEX OF A MONOMETAL METALLOCARBORANE C-ETC(U)


MAR 82 W C KALB, Z DENIDOWICZ, D W SPECKMAN

N00014-76-C-0390

TR-116

NL

UNCLASSIFIED

for 1  
ADVIS


END

DATE

FWD

BY

NTIC



AD A11 3192

12

OFFICE OF NAVAL RESEARCH

Contract No. N00014-76-C-0390

Task No. NR 053-608

TECHNICAL REPORT NO. 116

REACTIONS AT THE METAL VERTEX OF A MONOMETAL  
METALLOCARBORANE CLUSTER. THE CHEMISTRY OF  
[closo-3,3-(PPh<sub>3</sub>)<sub>2</sub>-3-(HSO<sub>4</sub>)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>]  
AND [closo-3-(PPh<sub>3</sub>)-3,3-(NO<sub>3</sub>)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>].

By

William C. Kalb, Zenon Demidowicz, Donna M. Speckman,  
Carolyn Knobler, Raymond G. Teller and M. Frederick  
Hawthorne\*

Prepared for Publication  
in  
Inorganic Chemistry

Department of Chemistry  
University of California  
Los Angeles, California 90024

March 9, 1982

Reproduction in whole or part is permitted for  
an purpose of the United States Government

Approved for Public Release; Distribution Unlimited

DTIC FILE COPY

DTIC  
FILED  
APR 8 1982  
H

00 04 08 030

REACTIONS AT THE METAL VERTEX OF A MONOMETAL  
METALLOCARBORANE CLUSTER. THE CHEMISTRY OF  
[closo-3,3-(PPh<sub>3</sub>)<sub>2</sub>-3-(HSO<sub>4</sub>)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>]  
AND [closo-3-(PPh<sub>3</sub>)-3,3-(NO<sub>3</sub>)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>].

By

William C. Kalb, Zenon Demidowicz, Donna M. Speckman, Carolyn Knobler,  
Raymond G. Teller and M. Frederick Hawthorne\*

Department of Chemistry  
University of California  
Los Angeles, California 90024

# ABSTRACT

Reaction of [closo-3,3-(PPh<sub>3</sub>)<sub>2</sub>-3-(H)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>](1) with sulfuric or nitric acid affords [closo-3,3-(PPh<sub>3</sub>)<sub>2</sub>-3-(HSO<sub>4</sub>)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>](2) or [closo-3-(PPh<sub>3</sub>)-3,3-(NO<sub>3</sub>)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>](3), respectively. Compound (3) can also be prepared from nitric acid and the dimeric metallocarborane [closo-Rh(PPh<sub>3</sub>)(C<sub>2</sub>B<sub>9</sub>H<sub>11</sub>)]<sub>2</sub> or from NO<sub>2</sub>/N<sub>2</sub>O<sub>4</sub> and (1). Complexes (2) and (3) have been used to prepare other new metallocarboranes; namely, [closo-3-(PPh<sub>3</sub>)-3,3-{C(Ph)-C(PPh<sub>3</sub>)-C(H)-C(Ph)}-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>](5), [closo-3-(PPh<sub>3</sub>)-3-(μ-CN)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>]<sub>4</sub>(7), [closo-3-(PPh<sub>3</sub>)-3-(L)-3-(NO<sub>3</sub>)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub> (L = CO, (8); L = PPh<sub>3</sub>, (9)), [closo-3,3-(PMe<sub>2</sub>Ph)<sub>2</sub>-3-NO<sub>3</sub>-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>](10) and [closo-3-(PPh<sub>3</sub>)-3-(CO)-3-(Cl)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>](11). Complexes (5) and (7) have been characterized by X-ray crystallography. The reactions of these new metallocarboranes described herein are representative of interconversions carried out at a discrete transition metal vertex of a cluster species.

Complex (5) crystallizes in space group P1̄ with 2 formula units in a cell of dimensions  $a = 12.763(6) \text{ \AA}$ ,  $b = 13.348(5) \text{ \AA}$ ,  $c = 14.561(7) \text{ \AA}$ ,  $\alpha = 91.58(3)^\circ$ ,  $\beta = 93.72(3)^\circ$ , and  $\gamma = 74.64(3)^\circ$ . Data were collected at  $-154^\circ \text{ C}$  on a Picker FACS-1 diffractometer using the  $\theta$ - $2\theta$  scan method. Least-squares refinement, including anisotropic vibration parameters for Rh and P, isotropic vibration parameters for other nonhydrogen atoms, and with each phenyl group described as a rigid group having a single isotropic vibration parameter, led to final conventional agreement indices (on F) of  $R = 0.048$  and  $R_w = 0.051$ , based on 4493 unique reflections having  $I > 3\sigma(I)$ . The molecule consists of a [C<sub>2</sub>B<sub>9</sub>H<sub>11</sub>]<sup>2-</sup> cage and a triphenylphosphine ligand bound to the metal atom of the trisubstituted metallopentacycle Rh-C(Ph)-C(PPh<sub>3</sub>)-C(H)-C(Ph). Rh-C, Rh-B, B-B, B-C, and C-C distances are normal for a 3,1,2-RhC<sub>2</sub>B<sub>9</sub> closo-rhodacarborane fragment and the pattern of short-long-short C-C bond lengths in the RhC<sub>4</sub> ring is reminiscent of

-1a-

a penta-substituted cis-butadiene.

Complex (7)  $\cdot 5C_6H_6$  crystallizes in space group  $P2_1/a$  with 4 formula units in a cell of dimensions  $a = 26.046(8) \text{ \AA}$ ,  $b = 15.626(3) \text{ \AA}$ ,  $c = 30.355(8) \text{ \AA}$ , and  $\beta = 106.71(2)^\circ$ . Data were collected at  $-154^\circ \text{ C}$  on a Syntex  $P1$  diffractometer using the  $\theta$ - $2\theta$  scan method. Least-squares refinement, including vibration parameters and rigid group assignments as described above, led to final conventional agreement indices (on F) of  $R = 0.063$  and  $R_w = 0.078$ , based on 9732 unique reflections have  $I > 3\sigma(I)$ . The molecules consist of four discrete closo-phosphino-rhodacarborane moieties joined together through their respective metal vertices by cyano ligand bridges. Each tetramer possesses a crystallographic 2-fold axis; the 2 non-crystallographically equivalent tetramers are very similar. The Rh-Rh separation is approximately  $5 \text{ \AA}$ . Bond distances within each icosahedral fragment are normal for such a closo-Rh(III)-metallo-carborane.



Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A	

## INTRODUCTION

One of the more interesting challenges of the contemporary chemistry of metallocarboranes derived from transition metals is the development of new reactions and species which make use of the unique reactivity of the transition metal vertex. One approach which we have employed utilizes closo-metallocarboranes which bear dissociable ligands, such as phosphines and carbon monoxide, on their transition metal vertices. The several reports<sup>1</sup> of metallocarboranes of this type which have appeared were mainly concerned with molecular structure and dynamics. In this paper we report the synthesis and reactions of two rhodacarborane species, a bisulfate complex [closo-3,3-(PPh<sub>3</sub>)<sub>2</sub>-3-(HSO<sub>4</sub>)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>](2)<sup>2</sup> and a bidentate nitrate [closo-3-(PPh<sub>3</sub>)-3,3-(NO<sub>3</sub>)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>](3)<sup>3</sup>. The molecular structures of (2) and (3) have been reported elsewhere<sup>2,3</sup>.

## RESULTS AND DISCUSSION

The reaction of [closo-3,3-(PPh<sub>3</sub>)<sub>2</sub>-3-(H)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>](1) with either sulfuric or nitric acid in dichloromethane proceeds to give the bisulfate (2) or the nitrate (3), respectively. Compound (2) precipitates directly from the reaction mixture and can be isolated as orange crystals by vigorous trituration with diethylether and (3) can be purified by column chromatography and obtained as red micro-crystals. The nitrate (3) can be prepared more conveniently by the reaction of nitric acid with the dimeric metallocarborane [{closo-Rh(PPh<sub>3</sub>)(C<sub>2</sub>B<sub>9</sub>H<sub>11</sub>)}<sub>2</sub>]<sup>4</sup>, also in dichloromethane. Attempts to prepare the corresponding complex [closo-3-(PPh<sub>3</sub>)-3,3-(HSO<sub>4</sub>)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>](4) from [{closo-Rh(PPh<sub>3</sub>)(C<sub>2</sub>B<sub>9</sub>H<sub>11</sub>)}<sub>2</sub>] and sulfuric acid, under similar conditions, were unsuccessful. We also note that (3) may be prepared by the reaction of an NO<sub>2</sub>/N<sub>2</sub>O<sub>4</sub> gas mixture with (1).

Closo-complexes (2) and (3) are formally Rh(III)-metallocarboranes, and it is noteworthy that they represent the first reported examples of metallocarboranes which contain metal-oxygen bonds. In addition, (3) constitutes a



rare example of a complex in which a bidentate nitrate ligand and a strong  $\pi$ -acceptor, such as triphenylphosphine, share the same coordination sphere. A small number of stable platinum nitrate complexes containing  $\pi$ -acceptor ligands have been reported.<sup>5</sup>

As expected, the reactivities of the monodentate bisulfate (2) and the bidentate nitrate (3) differ. We have found that (2) reacts with hydrogen gas at room temperature to yield (1) and sulfuric acid. Substitution of deuterium gas for hydrogen results in the formation of virtually pure [closo-3,3-(PPh<sub>3</sub>)<sub>2</sub>-3-(D)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>] provided that one equivalent of triphenylphosphine is initially present in the reaction mixture to suppress carborane terminal B-H/B-D exchange.<sup>6</sup> In contrast, the reaction of (3) with hydrogen gives [closo-Rh(PPh<sub>3</sub>)(C<sub>2</sub>B<sub>9</sub>H<sub>11</sub>)]<sub>2</sub><sup>4</sup>, [closo-3,3-(PPh<sub>3</sub>)<sub>2</sub>-3-(H)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>] (1) and another, as yet uncharacterized, product all identified by <sup>1</sup>H, <sup>31</sup>P and <sup>11</sup>B NMR spectroscopy.<sup>7</sup>

Several mechanisms can be proposed for the reaction of (2) with hydrogen. Four possibilities are shown in Scheme I. The reaction may proceed through the neutral species [closo-3-(PPh<sub>3</sub>)-3,3-HSO<sub>4</sub>-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>] (4) produced by dissociation of PPh<sub>3</sub> from (2) or through an ionic species such as [closo-3,3-(PPh<sub>3</sub>)<sub>2</sub>-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>]<sup>+</sup>[HSO<sub>4</sub><sup>-</sup>] (4a). Secondly, the hypothetical intermediates (4) and (4a) may each react with hydrogen by one of two pathways; heterolytic cleavage to give (1) and sulfuric acid, or through the oxidative addition of hydrogen to generate unprecedented formal Rh(V) seven-coordinate dihydrides which reductively eliminate H<sub>2</sub>SO<sub>4</sub> or lose a proton to give the observed products. Oddly enough, a similar and isoelectronic Ru (IV) dihydride [closo-3,3-(PPh<sub>3</sub>)<sub>2</sub>-3,3-(H)<sub>2</sub>-3,1,2-RuC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>] has been characterized.<sup>8</sup> In order to evaluate the mode of reactivity of (2) with hydrogen, the solution behavior of (2) in acetonitrile was examined by <sup>31</sup>P FTNMR and conductivity measurements. The <sup>31</sup>P{<sup>1</sup>H} FTNMR spectrum of (2) in acetonitrile at 230K



revealed two doublets at +46.9 ppm and +36.1 ppm plus a singlet at -1.5 ppm which corresponded to free triphenylphosphine. The lower-field doublet was found to be characteristic of a mono-(PPh<sub>3</sub>) rhodacarborane ( $J_{Rh-P}=144$  Hz) and the higher field doublet is due to a bis-(PPh<sub>3</sub>) species ( $J_{Rh-P}=124$  Hz). The nature of the mono-(PPh<sub>3</sub>) compound was determined directly. This compound was isolated from acetonitrile as a yellow solid, characterized, and found to contain a bis(acetonitrile) triphenylphosphine rhodacarborane cation, [closo-3,3-(NCCH<sub>3</sub>)<sub>2</sub>-3-(PPh<sub>3</sub>)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>]<sup>+</sup>[HSO<sub>4</sub>]<sup>-</sup> (2a). The bis-(PPh<sub>3</sub>) complex was not isolable. However, conductivity measurements performed on the acetonitrile solution at 25°C tentatively indicated that the nature of this bis-(PPh<sub>3</sub>) species was also cationic. A plot of molar conductivity versus (concentration)<sup>1/2</sup> was linear over a concentration range of 1.1 mM to 6.4 mM and is indicative of a solution of strong 1:1 electrolytes. Confirmation of the cationic nature of the bis-(PPh<sub>3</sub>) species (and that of (2a) as well) was obtained by the addition of a bisulfate anion source,  $\mu$ -nitrido-bis(triphenylphosphorous)(1+) bisulfate, (PPN)<sup>+</sup>(HSO<sub>4</sub>)<sup>-</sup>, to the acetonitrile solution of (2). Addition of six equivalents of (PPN)<sup>+</sup>(HSO<sub>4</sub>)<sup>-</sup> caused the appearance of a new doublet in the <sup>31</sup>P NMR spectrum at +32.4 ppm ( $J_{Rh-P}=127$  Hz) with a corresponding reduction in all of the signals observed prior to the addition of the excess bisulfate. These results imply that the following equilibria exist in solution:



In the absence of excess bisulfate ion, no detectable amounts of (2) exist in solution. However, addition of (PPN)<sup>+</sup>(HSO<sub>4</sub>)<sup>-</sup> displaces the position of the equilibrium in step (A) in favor of the neutral compound (2) (which appears as the new doublet in the <sup>31</sup>P NMR spectrum) and this consequently causes a reduction

in the amounts of (2a), the bis-(PPh<sub>3</sub>) species, and free PPh<sub>3</sub>. The most likely formulation for the bis-(PPh<sub>3</sub>) species is [closo-3,3-(PPh<sub>3</sub>)<sub>2</sub>-3-(NCCH<sub>3</sub>)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>]<sup>+</sup>HSO<sub>4</sub><sup>-</sup>. Support for a solvent-containing bis-(PPh<sub>3</sub>) compound was provided by the observation that a solution of (2) in acetonitrile at 25°C does not react with hydrogen gas; however, the reaction does proceed rapidly at elevated temperatures to give (1) in quantitative yields. Thus, it was postulated that although the species in solution are cationic, strongly bound acetonitrile ligand is present, coordinatively saturating the metal center. This ligand must be thermally labilized in order to allow a hydrogen molecule to attack. Since a simple doublet was observed in the <sup>31</sup>P{<sup>1</sup>H} NMR spectrum of the bis-(PPh<sub>3</sub>) species, both triphenylphosphines are in identical environments. This is probably due to rapid rotation of the metal vertex with respect to the carborane ligand in the cationic bis-(PPh<sub>3</sub>) acetonitrile complex.

Since these observations prove that in solution compound (2) exists as a mixture of solvent-containing cations, the mechanism for the reaction of (2) with hydrogen gas probably resembles the case outlined in Scheme I in which the formation of a coordinatively unsaturated rhodacarborane cation was the necessary initial step. In the mechanistic scheme for the reaction of (2) with hydrogen gas we intuitively favor the heterolytic cleavage of dihydrogen.

Having found that a cation derived from (2) reacts with hydrogen, we explored the possibility that a reaction would occur between (2) and a reactive organic substrate such as a terminal acetylene. A solution of (2) in tetrahydrofuran reacts rapidly with an excess of phenylacetylene at 45°C to afford a yellow complex (5) in excellent yield. The compound (5) can also be

#### Scheme I

obtained from (3) and excess phenylacetylene in the presence of an equimolar

portion of triphenylphosphine. Compound (5) was characterized by spectroscopy and an X-ray crystallographic study. The structure of (5) is illustrated in Fig. 1

Figure 1

and some selected distances and angles are reported in Table I.

Table I

The molecule consists of a  $[C_2B_9H_{11}]^{2-}$  cage and a triphenylphosphine ligand bound to the metal atom of the trisubstituted metallocycle  $Rh-C(Ph)-C(PPh_3)-C(H)-C(Ph)$ . The molecular formula of (5) can thus be written as  $[closo-3-(PPh_3)-3,3-\{C(Ph)-C(PPh_3)-C(H)-C(Ph)\}-3,1,2-RhC_2B_9H_{11}]$ . The Rh-C, Rh-B, C-C, C-B and B-B distances are normal for a 3,1,2- $RhC_2B_9$  closo-rhodacarborene fragment. The metallocycle is best represented as shown in Figure 2, with double bonds localized as indicated.

Figure 2

This description is supported by the following structural observations:

- The angle between the plane of the flat  $RhC_4$  ring and the respective plane of the two phenyl substituents ( $42^\circ$  and  $63^\circ$ ) rule out the significant  $\pi$ - $\pi$  overlap.
- The C(3)-phenyl and C(6)-phenyl bond lengths (1.509(8) and 1.513(8) Å respectively) shown no double bond character.
- The Rh-C(3) and Rh-C(6) bond distances (2.058(6) and 2.069(6) Å, respectively) are within the range found for Rh(III)-C single bonds.<sup>9-12</sup>
- The four P-C bonds of the C(5)-bonded triphenylphosphonium group are virtually identical.
- The pattern of the short-long-short C-C bond lengths in the  $RhC_4$  ring (1.351(8), 1.454(8) and 1.363(8) Å, respectively) is reminiscent of a penta-substituted cis-butadiene.<sup>13</sup>

The  $^1\text{H}$  NMR spectrum of (5) contained two carborane C-H resonances and a signal at 6.86  $\delta$  assigned as a vinylic proton. The  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum showed two inequivalent phosphorus nuclei,  $\text{P}_1$  and  $\text{P}_2$  coupled to each other ( $J_{\text{P}_1-\text{P}_2} = 7$  Hz).  $\text{P}_1$  (a doublet of doublets) appeared at +30.9 ppm which is typical of a triphenylphosphine ligand bound directly to the metal vertex of a rhodacarborane fragment ( $J_{\text{Rh}-\text{P}_1} = 129$  Hz). On the other hand,  $\text{P}_2$  which is not directly coordinated to the metal vertex (also a doublet of doublets) appeared at much higher field (+3.6 ppm) and exhibited a  $J_{\text{Rh}-\text{P}_2}$  value of 12 Hz. The sharp nature of the signal observed for  $\text{P}_2$  also is consistent with its location on a vinylic carbon.

A possible mechanism for the formation of (5) from (2) is shown in Scheme II.

#### Scheme II

The first step involves the dissociation of the bisulfate anion from (2) giving (4a). Next, a  $\sigma$ -acetylide complex is formed with the elimination of sulfuric acid and the subsequent loss of triphenylphosphine is followed by complexation of a second phenylacetylene molecule. Insertion of the coordinated phenylacetylene molecule into the Rh-C bond of the  $\sigma$ -phenylacetylide ligand may then take place. Coordination of an additional acetylene molecule and subsequent insertion into a metal-carbon bond has been observed in the linear oligomerization of acetylene catalyzed by  $[\text{Ni}(\text{CO})_2(\text{PPh}_3)_2]$ .<sup>14</sup> It is noteworthy that in the present case the insertion step occurs in such a fashion that the phenyl substituent is in a position alpha to the rhodium atom and that only one isomer is formed in the reaction. A similar mode of addition has been noted in the reaction of  $[\text{Ni}(\text{CH}_3)(\text{PPh}_3)(\text{acac})]$  with phenylacetylene<sup>15</sup>; the vinyl complex formed by the addition of the Ni-CH<sub>3</sub> moiety to the phenylacetylene triple bond bears the phenyl substituent on the carbon atom immediately adjacent to the nickel center. In the final step, nucleophilic

attack of triphenylphosphine on the coordinated organic ligand occurs producing (5). An example of such an attack on a coordinated unsaturated molecule is the reaction of  $[\text{Os}_3(\text{H})(\text{CO})_{10}(\text{CH}=\text{CH}_2)]$  with dimethylphenylphosphine to give  $[\text{Os}_3(\text{H})(\text{CO})_{10}(\text{CHCH}_2\text{P}^+\text{Me}_2\text{Ph})]^{16}$ . The reaction of phenylacetylene with either (2) or (3), the latter in the presence of triphenylphosphine, to give (5) constitutes the first example of metallocycle construction at a metal vertex of a metallocarborane cluster.

Not surprisingly, it was found that  $[\text{closo-3,3-(PPh}_3)_2\text{-3-(HSO}_4\text{)-3,1,2-RhC}_2\text{B}_9\text{H}_{11}](2)$  formally dehydrogenates simple alcohols to give the corresponding aldehydes. As examples one finds that (2) reacts with ethanol or *n*-propanol at slightly elevated temperatures to produce acetaldehyde or propionaldehyde, respectively. Complex (1) and sulfuric acid are also produced. This reaction occurs heterogeneously since both the reactant (2) and product (1) are insoluble in ethanol and *n*-propanol. Two possible mechanisms for this facile reaction are shown in Scheme III.

### Scheme III

In Path (1), loss of triphenylphosphine from (2) is followed by coordination of alcohol and in Path (2) loss of the bisulfate anion occurs before the alcohol becomes bound. Hydrogen transfer subsequent to this results in the elimination of the aldehyde from the alkoxyrhodacarborane species; formation of complex (1) in Path (1) requires the presence of triphenylphosphine.

In order to confirm that the hydroxy proton on the alcohol is indeed lost to form sulfuric acid and that an  $\alpha$ -hydrogen from the alcohol becomes the rhodium-bound hydride in complex (1) upon elimination of aldehyde, (2) was reacted with  $d_1$ -ethanol and with  $d_5$ -ethanol. The rhodacarborane products formed from these two reactions were examined by  $^1\text{H}$  NMR and by IR spectroscopy

to determine the amounts of Rh-H and Rh-D present. Compound (1) produced in the reaction of (2) with  $d_5$ -ethanol exhibited no Rh-H resonances in the  $^1\text{H}$  NMR and no Rh-H peaks in the infrared spectra. A weak band at  $1520\text{ cm}^{-1}$ , corresponding to a Rh-D stretch, was observed. The product (1) formed in the reaction of (2) with  $d_1$ -ethanol revealed a strong six-line pattern in the hydride region of the  $^1\text{H}$  NMR ( $-8.3\text{ ppm}$ ). A weak band was observed for the Rh-H stretch at  $2065\text{ cm}^{-1}$  in the infrared spectrum as well whereas no IR bands due to Rh-D could be detected. These results indicate that the fate of deuterium in O-D and C-D bonds is consistent with the two mechanisms presented. A mechanism similar to the one depicted in Path (1) of Scheme III has been proposed to account for the catalytic dehydrogenation of alcohols by nitrate and perfluorocarboxylate - ruthenium complexes.<sup>17</sup> In contrast, the reaction reported here proceeds stoichiometrically, and attempts to make it catalytic by the addition of sulfuric acid have been unsuccessful.

$[\text{closo-3,3-(PPh}_3)_2\text{-3-(HSO}_4\text{)-3,1,2-RhC}_2\text{B}_9\text{H}_{11}](2)$  reacts with sodium bromide (in benzene-water at reflux) or with hydrochloric acid (in tetrahydrofuran at room temperature) to afford, in good yield,  $[\text{closo-3,3-(PPh}_3)_2\text{-3-(X)-3,1,2-RhC}_2\text{B}_9\text{H}_{11}](6)(\text{X=Br or Cl})$ <sup>18</sup>. The bisulfate (2) also reacts with sodium cyanide to afford a yellow compound (7) initially formulated as  $[\text{closo-3,3-(PPh}_3)_2\text{-3-(CN)-3,1,2-RhC}_2\text{B}_9\text{H}_{11}]$ . The infrared spectrum of (7) contained bands typical of coordinated triphenylphosphine and carborane ligands. A strong, sharp peak at  $2146\text{ cm}^{-1}$  was also observed. This was attributed to  $\nu(\text{CN})$  and seemed characteristic of a terminal cyano ligand coordinated to a Rh(III) center.<sup>19</sup> In addition, the  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum of (7) showed one doublet at  $+40.9\text{ ppm}$  ( $J_{\text{Rh-P}}=117\text{ Hz}$ ). However, elemental analyses and molecular weight studies were consistent with the formulation of (7) as  $[\text{Rh}(\text{CN})(\text{PPh}_3)(\text{C}_2\text{B}_9\text{H}_{11})]_4$ . In order to determine the exact nature of (7) an X-ray crystallographic study was carried out. One of the two crystallographically unique

molecules of (7) is illustrated in Figure 3, and some distances and angles are shown in Table II.

Figure 3

Table II

The molecule has a cyclic structure consisting of four discrete closo-phosphino-rhodacarborane moieties joined together through their respective metal vertices by essentially linear cyano ligand bridges. The formula of (7) can thus be written as [closo-3-(PPh<sub>3</sub>)-3-(μ-CN)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>]<sub>4</sub>. Metal-metal bonds are absent in (7) and the Rh-Rh separation is approximately 5Å, the structure being held together entirely by the cyano groups. The bond distances within each RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub> fragment are normal for such a closo-Rh(III)-metallocarborane and, assuming that each carborane cage occupies three coordination sites, the geometry about the rhodium centers can be classified as octahedral. The triphenylphosphine ligands are unexceptional and are staggered with respect to one another, probably for steric reasons. Although the Rh-cyano group angles are all nearly linear, 170-176°, the torsion angles Rh-C-N-Rh range from 16° to 78°; the (RhCN)<sub>4</sub> moiety is not planar. At first, this may appear to be due to packing forces, but the fact that the two unique tetrameric molecules exhibit nearly identical angles of twist indicates that this distortion from planarity is due to electronic or intramolecular steric effects. Although dimeric and polymeric transition metal complexes containing bridging cyano ligands are well known<sup>20</sup> we believe that (7) represents the first example of a discrete tetramer with such linkages. Finally we note that, contrary to our expectation, the reaction of (3) with sodium cyanide under analogous conditions gave only an intractable mixture of products.

It is well known that the nitrate ligand may be either a bidentate four-electron donor or a monodentate two-electron donor towards a transition metal center.<sup>21</sup> Accordingly, we have studied the reaction of (3) with donor ligands

$L(L=CO, PPh_3 \text{ or } PMe_2Ph)$  in the hope that a complex of the type  $[closo-3-(PPh_3)-3-(L)-3-(NO_3)-3,1,2-RhC_2B_9H_{11}]$ , containing the relatively uncommon monodentate nitrate group, would be formed.

The reaction of (3) with carbon monoxide or triphenylphosphine in diethyl-ether gave good yields of  $[closo-3-(PPh_3)-3-(CO)-3-(NO_3)-3,1,2-RhC_2B_9H_{11}]$  (8) or  $[closo-3,3-(PPh_3)_2-3-(NO_3)-3,1,2-RhC_2B_9H_{11}]$  (9), respectively. However, reaction of (3) with excess dimethylphenylphosphine resulted in the displacement of triphenylphosphine with the formation of  $[closo-3,3-(PMe_2Ph)_2-3-(NO_3)-3,1,2-RhC_2B_9H_{11}]$  (10) in excellent yield. Complexes (8), (9) and (10), formally Rh(III)-metallo-carboranes containing monodentate  $NO_3$  ligands, were characterized by elemental analyses, infrared and NMR spectroscopy.

As previously reported<sup>3</sup> solutions of (9) exhibit  $^{31}P\{^1H\}$  NMR spectra which confirm the presence of uncoordinated triphenylphosphine. This observation is consistent with the following equilibrium.



We have also observed that (8) slowly loses its carbonyl ligand in solution. Thus the  $^{31}P\{^1H\}$  NMR spectrum of (8) initially contains an intense doublet at +41.2 ppm ( $J_{Rh-P}=107$  Hz) together with a weak resonance due to (3). Over a period of weeks the signal due to (8) disappears and is replaced by a strong resonance characteristic of (3). The  $^{31}P\{^1H\}$  NMR spectrum of (10) consists of only one doublet at +11.2 ppm ( $J_{Rh-P}=132$  Hz) and no uncoordinated dimethylphenylphosphine is apparent. Thus, in contrast to (2), (8) and (9) the species (10) shows no apparent tendency to dissociate its ligands in solution.

Like (2),  $[closo-3,3-(PPh_3)_2-3-(NO_3)-3,1,2-RhC_2B_9H_{11}]$  (9) reacts with hydrogen to afford  $[closo-3,3-(PPh_3)_2-3-(H)-3,1,2-RhC_2B_9H_{11}]$  (1). In contrast, (10) does not react with hydrogen in benzene under similar conditions. Finally, (10) reacts with hydrochloric acid to give the known compound



[closo-3,3-(PMe<sub>2</sub>Ph)<sub>2</sub>-3-(Cl)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>]<sup>22</sup> in quantitative yield.

The synthesis of the nitrate rhodacarboranes (3), (8), (9) and (10) offered a possible opportunity for distinguishing between mono- and bidentate nitrate ligands on the basis of their infrared absorption bands. However, on comparison of the infrared spectra of (3), (8), (9) and (10) (as nujol mulls) it became apparent that no simple distinction could be made with confidence. All four complexes exhibited a strong band in the regions 1515-1450 cm<sup>-1</sup> and 1282-1220 cm<sup>-1</sup>, no other obvious distinguishing features being observed. The presence of phosphine and carborane ligands also hampered the identification of any other lower frequency nitrate ligand vibrations which may have been present. In this context we note that in (3) the  $\nu(\text{NO}_3)$  band in the region 1282-1220 cm<sup>-1</sup> often appeared as two absorptions, closely spaced and of nearly equal intensity, apparently as a result of solid-state splitting.

The synthesis of [closo-3,3-(PPh<sub>3</sub>)<sub>2</sub>-3-(X)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>] (X=H, Br or Cl)<sup>18</sup> can be achieved easily and conveniently from (3) by direct reaction with triphenylphosphine and either hydrogen, tetraethylammonium bromide or hydrochloric acid. Similarly the complex [closo-3-(PPh<sub>3</sub>)-3-(CO)-3-(Cl)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>] (11) can be prepared directly from (3) by reaction with carbon monoxide followed by addition of hydrochloric acid. Compound (11) was characterized by elemental analyses, infrared and NMR spectroscopy. Previous attempts to prepare (11) by displacement of triphenylphosphine from (6) (X=Cl) with carbon monoxide had been unsuccessful.<sup>23</sup> In contrast to (8), (11) does not appear to lose its carbonyl ligand in solution. Some further facile reactions involving (6), (8), (9) and (11) are illustrated in Scheme IV.

#### Scheme IV

Finally, we note that the complexes [closo-3-(PPh<sub>3</sub>)-3,3-{C(Ph)-C(PPh<sub>3</sub>)-C(H)-C(Ph)}-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>] (5), [closo-3-(PPh<sub>3</sub>)-3-(CO)-3-(NO<sub>3</sub>)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>] (8) and [closo-3-(PPh<sub>3</sub>)-3-(CO)-3-(Cl)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>] (11) are racemic and chiral at their metal vertices.

#### EXPERIMENTAL SECTION

In general, manipulations were carried out in an inert atmosphere using standard Schlenk techniques or on a high vacuum line. Unless otherwise specified all solvents were purified by standard procedures and distilled under argon before use. Ethanol (absolute or 95%) was used without further treatment. The reagents employed were obtained commercially and were at least of reagent grade quality. NO<sub>2</sub>/N<sub>2</sub>O<sub>4</sub> mixture, hydrogen, deuterium (99.5 atom % D) and carbon monoxide were obtained from Liquid Carbonic and used without further treatment. Phenylacetylene (Aldrich) was distilled under vacuum before use and n-propanol (Aldrich) was distilled from anhydrous potassium carbonate. [closo-3,3-(PPh<sub>3</sub>)<sub>2</sub>-3-(H)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>] (1)<sup>1a</sup> and [closo-Rh(PPh<sub>3</sub>)(C<sub>2</sub>B<sub>9</sub>H<sub>11</sub>)<sub>2</sub>]<sup>4</sup> were prepared by literature methods.

Infrared spectra were obtained as nujol mulls on a Perkin-Elmer 137 or 521 spectrometer and were calibrated against the 1601 cm<sup>-1</sup> band of polystyrene. <sup>31</sup>P, <sup>11</sup>B and <sup>1</sup>H NMR spectra were obtained on a Bruker WP200 FT spectrometer. Additional <sup>1</sup>H NMR spectra were measured on a Varian Associates A-60 instrument and other <sup>11</sup>B NMR spectra were recorded on a spectrometer built by Professor F. Anet of the Department of Chemistry, University of California, Los Angeles. <sup>1</sup>H NMR spectra are with reference to internal tetramethylsilane; <sup>31</sup>P NMR spectra are calibrated against an external D<sub>3</sub>PO<sub>4</sub> standard and <sup>11</sup>B NMR spectra are with reference to external BF<sub>3</sub>·OEt<sub>2</sub>; all chemical shifts (reported in ppm) are quoted as positive downfield. Conductimetric measurements were performed using a Beckman Conductivity Bridge, Model RC-19 and platinum electrodes.

Analyses and molecular weight determinations were by Schwarzkopf Microanalytical Laboratories, New York.

Preparation of [closo-3,3-(PPh<sub>3</sub>)<sub>2</sub>-3-(HSO<sub>4</sub>)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>] (2)

Sulfuric acid (95-98%, 3.0 ml, 50 mmol.) was added to a solution of (1) (2.0g, 2.62 mmol.) in dichloromethane (100 ml) in the air and the mixture was stirred rapidly for 15 minutes. The resulting gelatinous precipitate was filtered off, washed with dichloromethane (3 X 20 ml) and then vigorously triturated with diethylether to give air stable crystals of (2) (1.3g, 56%). An analytical sample, as a tetrahydrofuran (THF)-solvate, was obtained by recrystallization from THF/n-heptane.

Analysis: Found; C, 54.51; H, 5.71; B, 10.66; Rh, 11.29; P, 6.47; S, 3.42%. Calculated for C<sub>42</sub>H<sub>50</sub>B<sub>9</sub>O<sub>5</sub>P<sub>2</sub>SRh; C, 54.30; H, 5.42; B, 10.47; Rh, 11.08; P, 6.67; S, 3.45%.

Infrared Spectrum<sup>24</sup>: 3065(w), 2600(vs), 2545(vs) 2530(vs), 1595(w), 1575(w), 1480(vs), 1435(vs), 1300(w), 1240(w), 1160(vs), 1090(m), 1050(vs), 1025(w), 1005(w), 880(m), 745(vs), 700(vs) cm<sup>-1</sup>.

NMR Spectra<sup>25</sup>: <sup>31</sup>P{<sup>1</sup>H} (81.02 MHz, THF/Me<sub>2</sub>SO, r.t.) +33.4 (J<sub>Rh-P</sub> = 152 Hz); a singlet due to uncoordinated triphenylphosphine is also observed. Cooling to -32°C gives the following: +36.2 (d, J<sub>Rh-P</sub> = 166 Hz), +33.0 (d, J<sub>Rh-P</sub> = 157 Hz), +26.8 (d, J<sub>Rh-P</sub> = 132 Hz) and a signal due to uncoordinated triphenylphosphine. <sup>11</sup>B{<sup>1</sup>H} (80.5 MHz, Me<sub>2</sub>SO, r.t.) -9.0 (s,br) and +5.0 (s,br).

Preparation of [closo-3,3-(NCCH<sub>3</sub>)<sub>2</sub>-3-(PPh<sub>3</sub>)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>]<sup>+</sup>[HSO<sub>4</sub>]<sup>-</sup> (2a).

A solution of (2) (1.35g, 1.45 mmol.) in acetonitrile (80 ml) was stirred for 10 minutes, the volume was reduced to 20 ml in vacuo, and 80 ml of diethylether was added with stirring. The resultant yellow precipitate was isolated, washed with 2 x 50 ml of diethylether and dried in vacuo to give (2a) (0.54g, 55%).

Analysis: Found; C, 42.43; H, 5.04; B, 13.90; Rh, 14.88; N, 4.23; P, 4.59; S, 4.89. Calculated for  $C_{24}B_9H_{33}RhN_2PSO_4$ ; C, 42.59; H, 4.92; B, 14.38; Rh, 15.21; N, 4.14; P, 4.58; S, 4.74.

Infrared Spectrum: 3350(w), 2600(s,sh), 2545(vs), 2500(s,sh), 2300(vw), 2280(vw), 1585(vw), 1565(vw), 1480(m), 1435(vs), 1320(w), 1240(vs), 1170(w), 1160(vs), 1120(vw), 1090(s), 1050(vs), 1020(w), 1005(w), 995(w), 900(vw), 855(m), 845(s), 750(vs), 700(vs).

NMR Spectra:  $^{31}P\{^1H\}$  (81.02 MHz,  $CH_3CN/CD_3CN$ , 230K) +46.0 (d,  $J_{Rh-P} = 144$  Hz)  $^{11}B\{^1H\}$  (126.7 MHz,  $CH_3CN$ , r.t.) -23.4 (s,br), -7.6 (s,br), -3.9 (s,br), +11.0 (s,br), and +13.1 (s,br).

Preparation of [closo-3-(PPh<sub>3</sub>)-3,3-(NO<sub>3</sub>)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>] (3).

- a) A solution of (1) (2.92g, 3.83 mmol.) in reagent grade (RG) dichloromethane (200 ml) in the air was treated with nitric acid (70%, 0.5 ml, 7.9 mmol) and the resulting red mixture stirred at room temperature for 30 minutes. The solution was then evaporated to dryness and the residue chromatographed on a silica gel column prepared in dichloromethane (RG). Elution with dichloromethane gave a broad red band turning blue at the front. This was collected and treated with nitric acid (70%, 10 drops from a dropping pipette) with stirring for 10 minutes. Addition of *n*-heptane (200 ml, RG) followed by removal of dichloromethane in vacuo gave a red microcrystalline precipitate of (3) (1.55g, 72%).
- b) To a solution of [closo-Rh(PPh<sub>3</sub>)(C<sub>2</sub>B<sub>9</sub>H<sub>11</sub>)]<sub>2</sub><sup>4</sup> (1.0g, 1.01 mmol) in dichloromethane (200 ml RG) under argon was added nitric acid (70%, 0.5 ml, 7.9 mmol) and the mixture stirred for 10 minutes at room temperature. Addition of *n*-heptane (100 ml, RG) to the bright-red solution followed by removal of dichloromethane in vacuo gave air stable red microcrystalline (3) (1.08g, 96%).

c) A gentle stream of  $\text{NO}_2/\text{N}_2\text{O}_4$  mixture was bubbled through a stirred yellow suspension of (1) (0.51 g, 0.67 mmol) in benzene (130 ml) under argon at room temperature. After 15 minutes the gas mixture source was removed, stirring continued for a further 5 minutes after which time the dark red solution was evaporated to dryness in vacuo. The residue was chromatographed on a silica gel column prepared in hexane (RG) under argon. Elution with dichloromethane (RG) gave a red band which was collected and treated with hexane (10 ml, RG). Evaporation and further hexane addition (10 ml) gave, on standing overnight, red needle crystals of (3) (0.13 g, 34%).

Infrared Spectrum: 3040(s), 2604(s,sh), 2571(s,sh) 2538(vs), 1590(vw), 1565(m), 1506(vs), 1473(s), 1431(s), 1326(w), 1309(w), 1252(s), 1236(s), 1221(m), 1192(w), 1164(w), 1144(w), 1094(s), 1076(w), 1065(vw), 1032(w), 1011(s), 1001(m), 981(m), 936(vw), 923(vw), 902(w), 870(vw), 859(vw), 848(vw), 801(vw), 791(w), 755(s), 750(s,sh), 728(m,sh), 712(s), 697(vs)  $\text{cm}^{-1}$ .

NMR Spectra:  $^{31}\text{P}\{^1\text{H}\}$  (81.02 MHz,  $\text{CDCl}_3$ , r.t.) +36.1(d,  $J_{\text{Rh-P}}=169$  Hz).  $^{11}\text{B}\{^1\text{H}\}$  (64.15 MHz,  $\text{CDCl}_3$ , r.t.) -25.7(s,br), -9.2(s,br), -3.1(s,br) and +11.8(s,br).

Reaction of (2) with Hydrogen or Deuterium.

A stirred solution of (2) (0.2 g, 0.22 mmol.) in THF (100 ml) was treated with hydrogen gas (1 atmosphere) for 10 minutes at room temperature. Addition of water (100 ml) followed by removal of THF in vacuo gave yellow microcrystalline (1) (0.15 g, 82%). A similar reaction of (2) with deuterium in the presence of one equivalent of triphenylphosphine gave [closo-3,3-( $\text{PPh}_3$ )<sub>2</sub>-3-(D)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>] in 88% yield (infrared spectrum shows  $\nu(\text{Rh-D})$  at  $1520\text{cm}^{-1}$ , nujol mull).

Preparation of [closo-3-( $\text{PPh}_3$ )-3,3-{C(Ph)-C( $\text{PPh}_3$ )-C(H)-C(Ph)}-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>] (5).

a) To a slurry of (2) (0.41 g, 0.44 mmol.) in THF (20 ml) under nitrogen was added phenylacetylene (2.0 ml, 20 mmol.) with stirring. The mixture was then

warmed to 45° and after 2 minutes a yellow precipitate had formed. After 20 minutes the mixture was cooled to room temperature, the precipitate was isolated, washed with *n*-hexane (3 x 10 ml), and dried in vacuo giving (5) (0.41 g, 97%). An analytical sample was obtained by recrystallization from dichloromethane/ethanol.

Analysis: Found; C, 66.47; H, 5.52; B, 9.32; Rh, 10.02; P, 6.30%. Calculated for  $C_{54}H_{52}B_9P_2Rh$ ; C, 66.37; H, 5.44; B, 10.10; Rh, 10.68; P, 6.43%.

Infrared Spectrum: 3050(w), 2560(vs), 2520(vs), 1590(m), 1575(w), 1460(vs), 1435(vs), 1375(vs), 1315(w), 1195(w), 1160(w), 1105(vs), 1090(vs), 1030(w), 1005(m), 990(w), 791(w), 760(vs), 720(vs), 700(vs)  $cm^{-1}$ .

NMR Spectra:  $^1H$  (200.12 MHz,  $CD_2Cl_2$ , r.t.) + 7.7 - 7.0 (m, 40, Ph), +6.86 (s, 1, vinyl C-H), +2.4 (s, 1, carborane C-H) and +2.1 (s, 1, carborane C-H).  $^{31}P\{^1H\}$  (81.02 MHz,  $d_5$  - pyridine, r.t.) +30.9 (d of d,  $J_{Rh-P_1}=129$  Hz,  $J_{P_1-P_2}=7$  Hz), + 3.6 (d of d,  $J_{Rh-P_2}=12$  Hz,  $J_{P_2-P_1}=7$  Hz).

b) Triphenylphosphine (0.15 g, 0.57 mmol.) was added at room temperature to a stirred solution of (3) (0.3 g, 0.54 mmol) in THF (150 ml) under argon. After 15 minutes phenylacetylene (1.0 ml, 10 mmol.) was added to the stirred solution. After 20 hours *n*-heptane (100 ml) was added to the now yellow-orange solution; removal of THF in vacuo gave a yellow precipitate which was filtered off in air, washed with petroleum ether (35-60°, RG) and dried giving (5) (0.51 g, 98%) as air stable yellow microcrystals identified by infrared and  $^{31}P$  NMR spectroscopy.

#### Reaction of (2) with Alcohols.

a) With ethanol: Complex (2) (0.08 g, 0.09 mmol.) was suspended in absolute ethanol (3.0 ml) and the mixture heated to 50° in a water bath with shaking. After 5 minutes a bright yellow crystalline precipitate had formed. After cooling the yellow product was filtered off, washed with ethanol (3 x 1 ml)

and diethylether (2 x 1 ml) and dried by suction giving (1) (0.58 g, 89%).

The reaction of (2) with  $d_1$ -ethanol was performed exactly as above to provide product (1) as well. The reaction of (2) with  $d_5$ -ethanol proceeded as follows: Complex (2) (0.09 g, 0.10 mmol) was suspended in a solution consisting of 1.27 ml (0.02 mol)  $d_6$ -ethanol and 0.8 ml (0.04 mol) distilled water. The mixture was heated as discussed above and the yellow product [closo-3,3-(PPh<sub>3</sub>)<sub>2</sub>-3-(D)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>] was isolated by filtration, rinsed with 3 ml of diethylether, and dried in vacuo. The production of acetaldehyde in the reaction was verified in a separate experiment as follows: Complex (2) (0.28 g, 0.30 mmol) was dissolved in *o*-dichlorobenzene (15 ml) under nitrogen. Absolute ethanol (5.0 ml) was then added with stirring. Nitrogen gas was continuously bubbled through the reaction mixture and the exit stream (with entrained acetaldehyde) was bubbled through a solution of 2,4-dinitrophenylhydrazine in sulfuric acid/ethanol/water. After 18 hours the indicator solution was extracted with benzene (4 x 10 ml) and the combined extracts were chromatographed on a basic alumina column prepared in benzene. Elution with dichloromethane gave a yellow band which was collected and evaporated to dryness in vacuo. This gave bright yellow crystals of acetaldehyde-2,4-dinitrophenylhydrazone (0.021 g, 31%, m.p. 157°; literature<sup>26</sup> m.p. = 157°).

b) With *n*-propanol: Complex (2) (0.25 g, 0.27 mmol) was suspended in *n*-propanol (10 ml) and the suspension heated to 50° in a water bath for 10 minutes. The resulting bright yellow crystalline precipitate was filtered off, washed with ethanol (3 x 4 ml) and diethylether (2 x 5 ml) and air-dried giving (1) (0.193 g, 94%).

The yield of propionaldehyde was determined in a separate experiment as follows: Complex (2) (0.3 g, 0.32 mmol) was charged into a 250 ml round-bottom flask equipped with a seal-off neck, a break-seal and a magnetic stir-bar. The flask was evacuated to a high vacuum through the seal-off neck.

n-Propanol (10 ml) was then distilled into the flask from anhydrous potassium carbonate (after four "freeze-pump-thaw" cycles to remove dissolved gases). The flask was then sealed under high vacuum and transferred to an oil bath at 75°. The suspension was stirred for 3 days. The flask was then cooled and the volatiles were removed by alembic distillation under high vacuum. The yield of propionaldehyde was determined to be 84% by g.l.c. analysis (4 ft. column, 10% Carbowax 20 M on Chromosorb P, acid washed).

Preparation of [closo-3,3-(PPh<sub>3</sub>)<sub>2</sub>-3-(Br)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>](6, X=Br)

a) Sodium bromide (1.2 g, 10 mmol) and complex (2) (0.5 g, 0.54 mmol) were heated together to the reflux temperature in benzene (150 ml) (to which water, 1 ml, had been added) for 30 minutes under nitrogen. The resulting mixture was cooled and filtered onto silica gel (20 ml dry volume). The resulting slurry was taken to dryness and applied to a silica gel column prepared in n-hexane. Elution with benzene produced a bright orange fraction which was collected and evaporated to dryness. Recrystallization of the residue from benzene/n-heptane gave (6, X=Br) as air stable orange crystals (0.29 g, 65%).

Analysis: Found; C, 54.60; H, 5.21; B, 11.60; Rh, 12.02; P, 7.34; Br, 8.94%.

Calculated for C<sub>38</sub>H<sub>41</sub>B<sub>9</sub>BrP<sub>2</sub>Rh; C, 54.35; H, 4.92; B, 11.58; Rh, 12.25; P, 7.38; Br, 9.52%.

Infrared Spectrum: 3000(w), 2540(vs), 2520(vs), 1590(w), 1570(w), 1480(vs), 1430(vs), 1265(w), 1185(w), 1160(w), 1100(vs), 1090(vs), 1080(s), 1025(w), 1020(w), 1005(m), 995(m), 895(w), 740(vs), 695(vs) cm<sup>-1</sup>.

NMR Spectra: <sup>1</sup>H (60 MHz, CD<sub>2</sub>Cl<sub>2</sub>, r.t.) + 7.5-7.2(m, 30, Ph), (s, 2, carborane C-H).

<sup>31</sup>P{<sup>1</sup>H} (81.02 MHz, CD<sub>2</sub>Cl<sub>2</sub>, r.t.) + 27.7 (d, J<sub>Rh-P</sub> = 134 Hz).

b) Complex (3) (0.4 g, 0.72 mmol) was dissolved in dichloromethane (100 ml, RG) at room temperature under argon and triphenylphosphine (0.24 g, 0.92 mmol) added with stirring. After 5 minutes tetraethylammonium bromide



(0.21 g, 1.0 mmol) was added and the mixture stirred for 45 minutes. The red solution was then chromatographed on a silica gel column prepared in dichloromethane (RG) in the air. Elution with dichloromethane gave a broad orange band which, when collected, treated with *n*-heptane (200 ml, RG) and evaporated, gave a red precipitate. This was washed with petroleum ether (20-40°, RG) and dried by suction giving (6) (X=Br) (0.59 g, approximately 98%) as a dichloromethane solvate identified by infrared spectroscopy.

Preparation of [*closo*-3-(PPh<sub>3</sub>)-3-(μ-CN)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>]<sub>4</sub> (7).

A mixture of (2) (1.07 mmol) and sodium cyanide (1.0 g, 20 mmol) in benzene (100 ml) and water (1.0 ml) was heated to reflux temperature with stirring under nitrogen for 20 minutes after which time the mixture was cooled to room temperature and filtered to remove excess cyanide (CARE!). The bright yellow filtrate was dried over anhydrous magnesium sulfate, filtered onto silica gel (10 ml dry volume) and the resulting slurry taken to dryness. The mixture was then applied to a silica gel column prepared in *n*-hexane. Elution with benzene gave a bright yellow band which, upon removal of the solvent, gave air stable (7) (0.46 g, 82%). An analytical sample was obtained by recrystallization from dichloromethane/ethanol.

Analysis: Found; C, 48.77; H, 5.38; B, 18.42; Rh, 19.12; P, 5.97; N, 2.55%. Calculated for C<sub>84</sub>H<sub>104</sub>B<sub>36</sub>N<sub>4</sub>P<sub>4</sub>Rh<sub>4</sub>; C, 48.17; H, 5.00; B, 18.58; Rh, 19.65; P, 5.91; N, 2.67%.

Molecular Weight: Found, 2043 (CHCl<sub>3</sub>). Calculated for C<sub>84</sub>H<sub>104</sub>B<sub>36</sub>N<sub>4</sub>P<sub>4</sub>Rh<sub>4</sub>; 2092.

Infrared Spectrum: 3048(w), 2570(vs), 2146(vs), 1595(w), 1580(w), 1480(vs), 1435(vs), 1200(w), 1170(w), 1100(vs), 1025(w), 1010(w), 990(m), 745(vs), 695(vs) cm<sup>-1</sup>.

NMR Spectra: <sup>1</sup>H(200.12 MHz, CD<sub>2</sub>Cl<sub>2</sub>, r.t.) + 7.6-7.3 (m, 15, Ph), + 3.2 (s, 1, carborane C-H), + 2.1 (s, 1, carborane C-H). <sup>31</sup>P{<sup>1</sup>H}(81.02 MHz, CD<sub>2</sub>Cl<sub>2</sub>, r.t.)

+40.9 (d,  $J_{\text{Rh-P}} = 117$  Hz).  $^{11}\text{B}\{^1\text{H}\}$  (111.8 MHz,  $\text{CHCl}_3$ , r.t.) + 8.5 (s,br), - 1.6 (s,br), - 5.3 (s,br) and - 8.8 (s,br).

Preparation of [closo-3-( $\text{PPh}_3$ )-3-(CO)-3-( $\text{NO}_3$ )-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>] (8).

A solution of (3) (0.174 g, 0.31 mmol.) in diethylether (50 ml) under argon was treated with carbon monoxide (1 atmosphere) with stirring at room temperature for 1-2 minutes (until the solution turned yellow from orange-red). *n*-Heptane (2 x 200 ml) was then added with rapid stirring. Cooling to 0° overnight gave a yellow precipitate which was filtered off in air, washed with *n*-pentane (RG) and dried by suction giving air stable (8) (0.13 g, 72%). An analytical sample was obtained by drying in vacuo.

Analysis: Found; C, 42.46; H, 4.60; N, 2.59; P, 5.36; B, 16.06; Rh, 16.44%. Calculated for C<sub>21</sub>H<sub>26</sub>B<sub>9</sub>NO<sub>4</sub>PRh; C, 42.92; H, 4.46; N, 2.38; P, 5.27; B, 16.56; Rh, 17.51%.

Molecular Weight: Found; 602 (toluene). Calculated for C<sub>21</sub>H<sub>26</sub>B<sub>9</sub>NO<sub>4</sub>PRh; 588.

Infrared Spectrum: 2994(m), 2538(vs), 2088(vs), 1656(vw), 1582(vw), 1565(vw), 1529(w), 1495(vs), 1473(m), 1431(s), 1326(vw), 1309(vw), 1260(vs), 1214(vw), 1192(vw), 1164(w), 1100(s), 1094(s,sh), 1076(vw), 1032(vw), 1016(w), 1001(m), 981(w), 967(vs), 940(vw), 923(vw), 898(vw), 859(vw), 852(vw), 804(w), 767(w,sh), 750(s), 725(m), 695(s) cm<sup>-1</sup>.

NMR Spectrum:  $^{31}\text{P}\{^1\text{H}\}$  (81.02 MHz,  $\text{CDCl}_3$ , r.t.) + 41.2 (d,  $J_{\text{Rh-P}} = 107$  Hz).

Preparation of [closo-3,3-( $\text{PPh}_3$ )<sub>2</sub>-3-( $\text{NO}_3$ )-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>] (9)<sup>27</sup>.

Triphenylphosphine (2.25 g, 0.95 mmol.) was added to a stirred solution of (3) (0.5 g, 0.9 mmol.) in diethylether (100 ml) under argon at room temperature and the resulting mixture stirred for 15 minutes. Addition of *n*-heptane (100 ml) and removal of diethylether in vacuo afforded an orange precipitate which was collected in air, washed with *n*-heptane and dried by suction giving air-stable (9) (0.59 g, 81%). The analytical sample was obtained by drying

in vacuo.

Analysis: Found; C, 55.62; H, 5.26; N, 1.58; P, 7.39; B, 11.62; Rh, 12.25%.

Calculated for  $C_{38}H_{41}B_9NO_3P_2Rh$ ; C, 55.53; H, 5.03; N, 1.70; P, 7.54; B, 11.84; Rh, 12.52%.

Infrared Spectrum: 3040(w), 2571(m), 2538(m), 1590(vw), 1577(vw), 1484(s), 1473(s,sh), 1431(s), 1318(vw), 1267(vs), 1192(w), 1164(w), 1088(m), 1032(vw,sh), 1022(w), 1006(m,sh), 991(m), 981(m,sh), 931(vw), 898(vw), 859(vw), 801(vw), 753(s), 700(s)  $cm^{-1}$ .

NMR Spectrum:  $^{31}P\{^1H\}$  (81.02 MHz,  $CDCl_3$ , r.t.) + 28.0 (d,  $J_{Rh-P}=134$  Hz); also observed are signals due to (3) and uncoordinated triphenylphosphine.

Preparation of  $[closo-3,3-(PMe_2Ph)_2-3-(NO_3)-3,1,2-RhC_2B_9H_{11}](10)$ .

Dimethylphenylphosphine (approximately 0.15 ml) was added to a stirred red solution of (3) (0.3 g, 0.54 mmol.) in diethylether (100 ml) at room temperature under argon. The solution turned yellow instantly. After 15 minutes *n*-heptane (50 ml) was added and removal of diethylether in vacuo gave a yellow crystalline precipitate. This was filtered off in air, washed with *n*-heptane (3 x 30 ml, RG) and dried by suction giving air-stable (10) 0.29 g, 94%). An analytical sample was obtained by drying in vacuo.

Analysis: Found; C, 37.38; H, 5.66; N, 2.50; P, 10.74; B, 16.65; Rh, 17.49%.

Calculated for  $C_{18}H_{33}B_9NO_3P_2Rh$ ; C, 37.69; H, 5.80; N, 2.44; P, 10.80; B, 16.96; Rh, 17.94%.

Molecular Weight: Found; 567 and 576 (benzene). Calculated for  $C_{18}H_{33}B_9NO_3P_2Rh$ ; 574.

Infrared Spectrum: 2994(w), 2538(vs), 1590(w), 1577(w), 1473(vs), 1462(m), 1431(s), 1326(m), 1318(m), 1300(m), 1284(s), 1260(vs), 1199(m), 1171(w), 1131(w), 1112(m,sh), 1100(s), 1082(w), 1065(w), 1022(s), 1006(s,sh), 986(vs), 953(vs),

927(vw), 918(vs), 910(vs), 886(w), 874(w), 859(m), 852(m), 804(m), 750(vs), 733(s), 715(s), 697(s)  $\text{cm}^{-1}$ .

NMR Spectra:  $^{31}\text{P}\{^1\text{H}\}$  (81.02 MHz,  $\text{CDCl}_3$ , r.t.) + 11.2 (d,  $J_{\text{Rh-P}}=132$  Hz).  $^{11}\text{B}\{^1\text{H}\}$  (127.04 MHz,  $\text{CHCl}_3$ , r.t.) -19.6(s), -15.5(s), -5.6(s), -2.4(s), +8.0(s).

Preparation of [closo-3-( $\text{PPh}_3$ )-3-(CO)-3-(Cl)-3,1,2-RhC $_2$ B $_9$ H $_{11}$ ](11).

a) Complex (3) (0.4 g, 0.72 mmol.) was dissolved in THF (100 ml under argon at room temperature) and carbon monoxide was passed through the stirred red solution until it turned yellow (about 1-2 minutes). Hydrochloric acid (37%, 0.2 ml, 2.4 mmol.) was then added dropwise and the resulting orange-yellow solution stirred for 15 minutes. Addition of *n*-heptane (100 ml), removal of THF in vacuo and further addition of *n*-heptane (70 ml) gave a golden yellow precipitate. This was filtered off in air, washed with *n*-heptane (RG) and dried by suction giving air-stable (11) (0.34 g, 85%).

Drying in vacuo gave an analytical sample.

Analysis: Found; C, 45.81; H, 4.63; Cl, 6.73; B, 17.58; P, 5.57; Rh, 18.20%.

Calculated for C $_{21}$ H $_{26}$ B $_9$ Cl OPRh; C, 44.96; H, 4.67; Cl, 6.32; B, 17.34; P, 5.52; Rh, 18.34%.

Molecular Weight: Found; 552 (benzene). Calculated for C $_{21}$ H $_{26}$ B $_9$ Cl OPRh; 561.

Infrared Spectrum: 2994(w), 2639(w), 2571(m), 2538(s), 2506(m), 2088(vs), 1590(w), 1577(w), 1484(m), 1431(s), 1318(w), 1199(w), 1164(w), 1125(vw), 1100(s), 1076(w), 1032(vw), 1016(w), 1006(w), 986(m), 940(vw), 931(vw), 923(vw), 859(vw), 764(m), 750(m), 744(m,sh), 725(w), 712(m), 696(s)  $\text{cm}^{-1}$ .

NMR Spectrum:  $^{31}\text{P}\{^1\text{H}\}$  (81.02 MHz,  $\text{CDCl}_3$ , r.t.) + 39.6 (d,  $J_{\text{Rh-P}}=107$  Hz).

b) A solution of (8) (0.14 g, 0.24 mmol.) in diethylether (100 ml), stirred at room temperature under argon, was treated with hydrochloric acid (37%, 3 drops from a dropping pipette). After 15 minutes *n*-heptane (200 ml) was added and the diethylether removed in vacuo. Further addition of *n*-heptane (200 ml) completed the precipitation of the product which was filtered off in

air, washed with *n*-heptane (RG) and dried by suction giving (11) (0.05 g, 20%) identified by infrared spectroscopy.

Formation of [closo-3,3-(PPh<sub>3</sub>)<sub>2</sub>-3-(H)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>](1)<sup>1a</sup> from (3) or (9) and Hydrogen.

- a) A solid mixture of (3) (0.09 g, 0.16 mmol) and triphenylphosphine (0.046 g, 0.18 mmol) was dissolved in THF (30 ml) under argon. Hydrogen gas (1 atmosphere) was then bubbled gently through the orange-red solution with stirring at room temperature for 10 minutes. The resulting yellow solution was treated with absolute ethanol (100 ml); evaporation in vacuo gave a yellow precipitate which was filtered off in air, washed with ethanol and diethylether, and dried by suction giving (1) (0.11 g, 90%) identified by infrared spectroscopy.
- b) Hydrogen (1 atmosphere) was bubbled through a stirred orange-red solution of (1) (0.2 g, 0.24 mmol) in THF (50 ml) at room temperature under argon. After 10 minutes absolute ethanol (150 ml) was added. Evaporation in vacuo gave a yellow precipitate which was collected in air, washed with ethanol and diethylether, and dried by suction giving (1) (0.15 g, 79%) identified by infrared spectroscopy.

Reaction of [closo-3-(PPh<sub>3</sub>)-3-(CO)-3-(NO<sub>3</sub>)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>] (8) with Triphenylphosphine.

A solid mixture of (8) (0.063 g, 0.11 mmol) and triphenylphosphine (0.063 g, 0.24 mmol) was dissolved in diethylether (70 ml) and the resulting solution stirred under argon at room temperature for 30 minutes. *n*-Heptane (200 ml) was then added and the diethylether removed in vacuo. Evaporation to a low volume gave an orange precipitate which was filtered off in air, washed with *n*-pentane (RG), and dried by suction giving (9) (0.068 g, 77%) identified by infrared spectroscopy.

Preparation of  $[\text{closo-3,3-(PPh}_3)_2\text{-3-(Cl)-3,1,2-RhC}_2\text{B}_9\text{H}_{11}]^{18}$  (6, X = Cl).

- a) A solid mixture of (3) (0.1 g, 0.18 mmol) and triphenylphosphine (0.05, 0.19 mmol) was dissolved in THF (30 ml) under argon. To the resulting solution was added, with stirring, hydrochloric acid (37%, 2 drops from a dropping pipette) and the mixture stirred for 10 minutes. Ethanol (100 ml) was then added and the mixture was reduced in volume by evaporation, giving a light-orange precipitate. This was collected in air, washed with ethanol and petroleum ether (35-60°, RG) and dried by suction giving (6, X = Cl) (0.12 g, 84%) identified by infrared spectroscopy.
- b) Four drops of hydrochloric acid (37%) were added to a stirred orange-red solution of (9) (0.2 g, 0.24 mmol) in THF (50 ml) at room temperature under argon. After 10 minutes ethanol (150 ml) was added. Evaporation to a low volume gave an orange precipitate which was filtered off in air, washed with ethanol and petroleum ether (35-60°, RG), and dried by suction giving (6, X = Cl) (0.17 g, 90%) identified by infrared spectroscopy.
- c) To a solution of (11) (0.3 g, 0.54 mmol) in THF (100 ml) under argon was added triphenylphosphine (0.15 g, 0.57 mmol) and the mixture stirred at room temperature for 15 minutes. Addition of *n*-heptane (50 ml) followed by evaporation gave an orange precipitate; further addition of *n*-heptane (50 ml) gave (6, X = Cl) (0.35 g, 81%) which was isolated in the air, washed with ethanol and *n*-pentane (RG), and dried by suction.
- d) A solution of (2) (0.3 g, 0.32 mmol) in THF (100 ml) under argon was treated with hydrochloric acid (37%, 6 drops from a dropping pipette) and the mixture stirred at room temperature for 15 minutes. Fifty milliliters of *n*-heptane was then added and THF removed in vacuo. The resulting light-orange precipitate was collected in air, washed with petroleum ether (35-60°, RG),

and dried by suction giving (6, X = Cl) (0.26 g, 93%) identified by infrared spectroscopy.

Reaction of [closo-3,3-(PMe<sub>2</sub>Ph)<sub>2</sub>-3-(NO<sub>3</sub>)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>] (10) with Hydrochloric Acid.

A yellow solution of (10) (0.1 g, 0.17 mmol.) in THF (100 ml) under argon was treated with hydrochloric acid (37%, 6 drops from a dropping pipette) and the mixture stirred under argon at room temperature. After 15 minutes a further 6 drops of acid were added. After 75 minutes *n*-heptane (100 ml) was added and the THF removed *in vacuo*. The yellow precipitate was filtered off in air, washed with *n*-pentane (RG) and dried by suction giving a compound (0.09 g) identified as [closo-3,3-(PMe<sub>2</sub>Ph)<sub>2</sub>-3-(Cl)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>]<sup>22</sup> (95%) by infrared spectroscopy.

X-ray Crystallographic Study of [closo-3-(PPh<sub>3</sub>)-3,3-{C(Ph)-C(PPh<sub>3</sub>)-C(H)-C(Ph)}-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>] (5) and [closo-3-(PPh<sub>3</sub>)-3-(μ-CN)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>]<sub>4</sub>·5C<sub>6</sub>H<sub>6</sub> (7).

Crystals of the two compounds chosen for data collection had the dimensions 0.10mm x 0.12mm x 0.10mm (5) and 0.20mm x 0.30mm x 0.45mm (7). They exhibited the faces (1 0 0), (1 0 0), (0 1 5), (0 1 5), (0 3 1), and (0 3 1) (5) and (6 1 0), (6 1 0), (1 0 3), (1 0 3), (0 1 0), and (0 1 0) (7), respectively. Data were collected and preliminary crystallographic experiments were performed on a Picker FACS-I diffractometer (5) and on a Syntex P1 diffractometer (7). Both crystals were maintained at -154°C by a cold nitrogen stream from an apparatus designed by Strouse.<sup>28</sup> Crystals of (7) from several solvent mixtures exhibited rapid deterioration, apparently due to solvent loss. Suitable crystals were finally produced from benzene/ethanol. After mounting the data-collection crystal of (7) it was immediately transferred to the cold nitrogen stream. The orientation of each crystal was determined by indexing reflections found photo-

graphically. For (7) an axial photograph revealed the presence of a mirror plane which suggested a monoclinic space group. For (5) the absence of symmetry suggested a triclinic cell. Preliminary data were collected to identify the intense high-angle data. Accurate cell constants were obtained by a least-squares fit of 15 (7) and 19 (5) high-angle reflections, respectively. Data were collected in the  $\theta$ - $2\theta$  scan mode with Zr-filtered  $\text{MoK}\alpha$  radiation for (5) and with graphite - monochromatized  $\text{MoK}\alpha$  radiation for (7). Three intense reflections were periodically monitored during data collection for each crystal. For (5) the intensities of the standards showed an isotropic decay of 8% and the raw data were corrected for this. Intensities of the standards for (7) fluctuated only slightly during the course of data collection. Other details of data collection are presented in Table V.

Table V

Data were corrected for Lorentz, polarization and absorption effects<sup>29</sup>. The data for (7) were corrected by an empirical method based on the fluctuation in intensities of 3 reflections near  $\chi = 90^\circ$ . The three reflections (and their  $2\theta$  values) are  $4\ 6\ \bar{2}$  ( $17^\circ$ ),  $7\ 9\ \bar{2}$  ( $26^\circ$ ), and  $9\ 11\ \bar{3}$  ( $32^\circ$ ). All three absorption curves exhibited the same shape with maxima and minima coinciding. The systematic absences  $h0l$ ,  $h=2n+1$  observed for (7) suggested the space groups  $\text{Pa}$  or  $\text{P2/a}$ . Successful refinement of the structure in the latter space group showed it to be the correct choice. An ignorance factor of 0.04 was used for each complex. For (5) only data satisfying the criteria  $I_o > 3\sigma(I_o)$  were retained for structure solution and refinement. For (7) only data satisfying the  $3\sigma$  criterion with  $\sin(\theta)/\lambda$  less than 0.40 were used in the structure analysis and only in the final cycles of refinement was the entire data set used.

The structures were solved by the heavy atom-Fourier synthesis method. In subsequent structure refinements the phenyl groups of the triphenylphosphine



moieties were treated as rigid groups with C-C and C-H bond distances set at 1.39 and 1.00 Å, respectively. During the least-squares process the positional and rotational parameters and the overall temperature factors of each group were allowed to vary and all the individual temperature factors were set at zero. For the non-group atoms all positional and thermal parameters (the Rh and P atoms were allowed to vibrate anisotropically, all other atoms isotropically) and the scale factor were varied in the full-matrix least-squares process.

A difference-Fourier synthesis of (5) after least-squares refinement to an agreement index of 0.054 revealed the eleven cage hydrogen atoms and the vinylic hydrogen of the metallocycle. The positional and thermal parameters of these atoms were also varied in three cycles of least-squares and convergence was realized at  $R(F) = 0.048$ . A final difference - Fourier synthesis showed no significant peaks.

For (7) all non-hydrogen atoms of the tetramer were varied in three cycles of least-squares refinement to an agreement factor 0.163. A difference - Fourier synthesis at this point showed peaks attributed to five molecules of benzene. Inclusion of these thirty atoms in the refinement lowered the agreement factor to 0.068. From this point on the full data set was used in the refinement (9732 observed reflections). Three cycles of full-matrix least-squares refinement converged to an agreement factor of 0.063. A final difference - Fourier synthesis revealed peaks attributable to hydrogen atoms on the cage atoms only. These atoms were not included in the model. All cyano group atoms were treated as half carbon, half nitrogen; an averaged carbon-nitrogen scattering factor was applied.

Final atomic parameters for (5) and (7) can be found in Table VI and VII.

Table VI

Table VII

REFERENCES

- 1) a) Paxson, T.E.; Hawthorne, M.F. J. Am. Chem. Soc. (1974), 96, 4674.
- b) Grimes, R.N. "Reactions of Metallocarboranes" in "Organometallic Reactions and Syntheses", Vol. 6, Plenum Press, New York, 1977, Chapter 2, pp. 63-221.
- c) Paxson, T.E.; Callahan, K.P.; Hoel, E.L.; and Hawthorne, M.F. "Metallocarboranes; Past, Present and Future" in Organotransition-Metal Chemistry", Y. Ishii and M. Tsutsui eds., Plenum Press, New York 1975, Chapter 1, pp. 1-23.
- d) Zakharkin, L.I.; L'vov, A.I. Zh. Obshch. Khim., (1971), 41, 1880.
- e) Hawthorne, M.F.; Ruhle, H.W. Inorg. Chem., (1969), 8, 176.
- f) Stone, F.G.A. J. Organometallic Chem., (1975), 100, 257.
- g) Colquhoun, H.M.; Greenhough, T.J.; Wallbridge, M.G.H. J.C.S. Chem. Commun., (1975) 1019; J. Chem. Soc. Dalton, (1978) 303.
- h) Colquhoun, H.M.; Greenhough, T.J.; Wallbridge, M.G.H. J.C.S. Chem. Commun., (1977) 737.
- i) Colquhoun, H.M.; Greenhough, T.J.; Wallbridge, M.G.H. J.C.S. Chem. Commun., (1978) 322.
- j) Miller, S.B.; Hawthorne, M.F. J.C.S. Chem. Commun., (1976) 786.
- k) Wong, E.H.; Hawthorne, M.F. J.C.S. Chem. Commun., (1976) 257; Inorg. Chem., (1978), 17, 2863.
- l) Teller, R.G.; Wilczynski, J.J.; Hawthorne, M.F. J.C.S. Chem. Commun., (1979) 472.
- m) Jung, C.W.; Hawthorne, M.F. J.C.S. Chem. Commun., (1976) 499; J. Amer. Chem. Soc., (1980), 102, 3024

- n) Jung, C.W.; Baker, R.T.; Knobler, C.B.; Hawthorne, M.F.  
J. Amer. Chem. Soc., (1980), 102, 5782.
- o) Kalb, W.C.; Kreimendahl, C.W.; Busby, D.C.; Hawthorne, M.F.  
Inorg. Chem., (1980), 19, 1590.
- p) Marder, T.B.; Long, J.A.; Hawthorne, M.F. J.C.S. Chem. Commun.,  
(1980), 677.
- q) Doi, J.A.; Teller, R.G.; Hawthorne, M.F. J.C.S. Chem. Commun., (1980), 80.
- r) Delaney, M.S.; Knobler, C.B.; Hawthorne, M.F. J.C.S. Chem. Commun., (1980),  
849; Inorg. Chem., Soc., (1981), 20, 1341.
- s) Jung, C.W.; Baker, R.T.; Hawthorne, M.F. J. Amer. Chem. Soc., (1981), 103, 810.
- t) Marder, T.B.; Baker, R.T.; Long, J.A.; Doi, J.A.; Hawthorne, M.F.  
J. Amer. Chem. Soc., (1981) 103, 2988.
- u) Delaney, M.S.; Teller, R.G.; Hawthorne, M.F. J.C.S. Chem. Commun., (1981) 235.
- v) Hewes, J.D.; Knobler, C.B.; Hawthorne, M.F. J.C.S. Chem. Commun., (1981) 206.
- 2) Kalb, W.C.; Teller, R.G.; Hawthorne, M.F. J. Amer. Chem. Soc., (1979), 101, 5417.
- 3) Demidowicz, Z.; Teller, R.G.; Hawthorne, M.F. J.C.S. Chem. Commun., (1979), 831.
- 4) Baker, R.T.; King, R.E., III; Knobler, C.; O'Con, C.A.; Hawthorne, M.F.  
J. Amer. Chem. Soc., (1978), 100, 8266.
- 5) Critchlow, P.B.; Robinson, S.D. Inorg. Chem., (1978), 17, 1896 and references  
therein.
- 6) Terminal B-H/B-D exchange with D<sub>2</sub> is catalyzed by complex (1); see E.L. Hoel  
and M.F. Hawthorne, J. Amer. Chem. Soc., (1974), 96, 4676; E.L. Hoel,  
M. Talebinasab-Savari and M.F. Hawthorne ibid. (1977), 99, 4356; W.C. Kalb,  
Ph.D. Thesis UCLA, 1979.
- 7) Baker, R.T.; Marder, T.B.; Hawthorne, M.F.; unpublished results.
- 8) Wong, E.H.; Hawthorne, M.F. Inorg. Chem., (1978), 17, 2863.

- 9) Mason, R.; Towl, A.D.C. J. Chem. Soc., (A) (1970) 1601.
- 10) Paulas, E.F.; Fritz, H.P; Schwarzhaus, K.E. J. Organometallic Chem., (1968), 11, 647.
- 11) Skapski, A.C.; Troughton, P.G.H. J.C.S. Chem. Commun., (1968), 525.
- 12) Skapski, A.C.; Troughton, P.G.H. ibid (1969), 666.
- 13) Cotton, F.A.; Wilkinson, G. "Advanced Inorganic Chemistry", 3rd edition, J. Wiley Interscience, New York. p. 732.
- 14) Meriwether, L.S.; Leto, M.F.; Colthup, E.C.; Kennerly, G.W. J. Org. Chem., (1962), 27, 3930.
- 15) Huggins, J.M.; Bergman, R.G. J. Amer. Chem. Soc., (1979), 101, 4410.
- 16) Churchill, M.R.; Shapley, J.R. J. Amer. Chem. Soc., (1976), 98, 2357.
- 17) Critchlow, P.B.; Robinson, S.D. Inorg. Chem., (1978), 17, 1902 and references therein.
- 18) a) Siedle, A.R. J. Organometallic Chem., (1975), 90, 249.  
b) Grey, R.; Hawthorne, M.F., unpublished results.
- 19) For comparison;  $[\text{Rh}(\text{H})_2(\text{CN})(\text{PPh}_3)_2]$ ,  $\nu(\text{CN})$  at  $2110 \text{ cm}^{-1}$ , Favero, G.; Rigo, P. Gazz. Chim. Ital., (1972), 102, 597.  $[\{\text{Rh}(\text{Cl})(\text{CN})_2\text{PPh}_3\}_2(\mu\text{-NCCN})]$ , terminal  $\nu(\text{CN})$  at  $2145$  and  $2135 \text{ cm}^{-1}$ , Favero, G.; Corain, B.; Rigo, P.; Turco, A. ibid (1973), 103, 297.
- 20) Sharpe, A.G; "The Chemistry of Cyano Complexes of the Transition Metals", Academic Press, New York 1976.
- 21) Critchlow, P.B.; Robinson, S.D. Coord. Chem. Rev., (1978), 25, 69.
- 22) Baker, R.T.; Ph.D. Thesis, UCLA, 1980.
- 23) Grey, R.; Hawthorne, M.F., unpublished results.
- 24) Intensities of the absorption bands are designated (in parentheses) as follows:  
vs, very strong; s, strong; m, medium; w, weak; vw, very weak; sh, shoulder.

- 25) s, singlet; d, doublet; m, multiplet; br, broad; r.t., room temperature.
- 26) "Handbook of Tables for Organic Compound Identification", p. 144, 3rd edition, CRC Press, Cleveland, Ohio 1967.
- 27) Difficulties were occasionally encountered in obtaining complex (9) by this method in a pure state, complex (3) appearing frequently as an impurity.
- 28) C.E. Strouse, Rev. Sci. Instrum., (1976), 47, 871.
- 29) Experimental and computational procedures used in this work have been previously described by K.P. Callahan, C.E. Strouse, A.L. Sims, and M.F. Hawthorne, Inorg. Chem., (1974), 13, 1397. The function minimized during least-squares refinement was  $\sum w(|F_o| - |F_c|)^2$  where  $w = 1/(\sigma F)^2$ .  $R = [\sum ||F_o| - |F_c|| / \sum |F_o|] / \sum |F_o|$ ;  $R_w = [\sum w||F_o| - |F_c||^2 / \sum w|F_o|^2]^{1/2}$ .

ACKNOWLEDGEMENTS

We thank C.A. O'Con, Dr. D. Busby and R.T. Baker for assistance in obtaining NMR spectra and Dr. R. Grey for preliminary experiments.

Financial support of this work by the National Science Foundation (Grant CHE78-05679) and the Office of Naval Research is gratefully acknowledged.

TABLE I

Some Distances (in Å) and Angles (in degrees) in [closo-3-(PPh<sub>3</sub>)-3,3-{C(Ph)-C(PPh<sub>3</sub>)-C(H)-C(Ph)}-3,1,2-RhC<sub>2</sub>[9H<sub>11</sub>](5).

## A. Distances

Rh-C(1)	2.279	Rh-B(8)	2.270
Rh-C(2)	2.284	Rh-C(3)	2.058
Rh-B(4)	2.251	Rh-C(6)	2.069
Rh-B(7)	2.328	Rh-P(1)	2.338
P(1)-Group 3	1.878	P(2)-Group 6	1.806
P(1)-Group 4	1.851	P(2)-Group 7	1.816
P(1)-Group 5	1.854	P(2)-Group 8	1.813
C(3)-Group 1	1.509	C(3)-C(4)	1.351
C(4)-HC4	0.98	C(4)-C(5)	1.454
C(5)-P(2)	1.795	C(5)-C(6)	1.363
C(6)-Group 2	1.513		
C(1)-C(2)	1.606	B(8)-B(9)	1.820
C(1)-B(4)	1.700	B(8)-B(12)	1.797
C(1)-B(5)	1.698	B(9)-B(10)	1.781
C(1)-B(6)	1.727	B(9)-B(12)	1.783
C(2)-B(6)	1.748	B(10)-B(11)	1.772
C(2)-B(7)	1.674	B(10)-B(12)	1.789
C(2)-B(11)	1.722	B(11)-B(12)	1.768
B(4)-B(5)	1.791	C(1)-H(1)	0.94
B(4)-B(8)	1.787	C(2)-H(2)	1.03
B(4)-B(9)	1.767	B(4)-H(4)	1.00
B(5)-B(6)	1.765	B(5)-H(5)	1.14
B(5)-B(9)	1.765	B(6)-H(6)	1.02
B(5)-B(10)	1.766	B(7)-H(7)	1.02
B(6)-B(10)	1.760	B(8)-H(8)	1.04
B(6)-B(11)	1.757	B(9)-H(9)	1.08
B(7)-B(8)	1.816	B(10)-H(10)	1.13
B(7)-B(11)	1.792	B(11)-H(11)	1.15
B(7)-B(12)	1.761	B(12)-H(12)	1.08

Table I (continued)

B. Angles

P(1)-Rh-C(3)	90.0	Cn <sup>a</sup> -Rh-C(3)	124.4
P(1)-Rh-C(6)	90.1	Cn <sup>a</sup> -Rh-C(6)	127.1
C(3)-Rh-C(6)	80.1	Cn <sup>a</sup> -Rh-P(1)	120.1
Rh-C(3)-C(4)	113.8	Rh-C(6)-C(5)	113.4
Rh-C(3)-Group 1	124.8	Rh-C(6)-Group 2	125.7
C(4)-C(3)-Group 1	121.9	C(5)-C(6)-Group 2	121.2
C(3)-C(4)-C(5)	116.7	C(6)-C(5)-C(4)	115.8
C(3)-C(4)-HC4	121.(3)	C(6)-C(5)-P(2)	128.2
C(5)-C(4)-HC4	121.(3)	C(4)-C(5)-P(2)	113.4

a) Centroid of the C<sub>2</sub>B<sub>3</sub> face of carborane.

Estimated standard deviations in bond lengths are: Rh-P 0.001, Rh-C 0.006, Rh-B 0.007, P-C 0.006, C-C 0.008, C-B 0.009, C-H 0.04, B-H 0.06; and angles are: Rh-C-C 0.4, P-Rh-C 0.2, P-Rh-Cn 0.2, C-Rh-C 0.2, C-C-C 0.3, C-C-P 0.4.



TABLE II

Some Distances<sup>a</sup> (in Å) and Angles (in degrees) in  
 $[\text{closo-3-(PPh}_3\text{)-3-(}\mu\text{-CN)-3,1,2-RhC}_2\text{B}_9\text{H}_{11}\text{]}_4\text{](7)}$ .

A. Distances	Monomer 1	Monomer 2	Monomer 3	Monomer 4
Rh-P	2.307(3)	2.322(3)	2.307(3)	2.298(3)
Rh-CN	2.031(9)	2.049(8)	2.057(8)	2.047(8)
	2.046(9)	2.054(8)	2.046(8)	2.060(9)
CN-CN <sup>b</sup>	1.162(16)	1.149(16)	1.119(12)	1.147(13)
CN-CN <sup>c</sup>	1.149(11)		1.136(12)	
Rh-C(carborane)	2.217(12)	2.260(11)	2.209(12)	2.242(11)
	2.219(11)	2.204(12)	2.245(12)	2.244(11)
Rh-B(carborane)	2.207(14)	2.238(12)	2.226(12)	2.217(11)
	2.222(12)	2.201(11)	2.243(12)	2.219(11)
	2.204(11)	2.197(12)	2.211(12)	2.199(11)
P-Phenyl 1 <sup>d</sup>	1.840	1.827	1.841	1.819
P-Phenyl 2	1.819	1.813	1.802	1.819
P-Phenyl 3	1.826	1.822	1.823	1.834
C-C(carborane)	1.655(15)	1.680(15)	1.625(15)	1.644(15)
C-B(carborane)	1.735(12)	1.730(31)	1.729(24)	1.741(15)
B-B(carborane)	1.780(4)	1.785(6)	1.793(4)	1.783(7)
B. Angles				
P-Rh-CN	90.5(3)	89.0(3)	94.6(3)	86.9(3)
	87.5(3)	95.9(3)	87.2(3)	91.9(3)
CN-Rh-CN	90.3(3)	86.9(3)	87.1(3)	89.2(3)
Rh-CN-CN	174.3(9)	170.5(9)	173.4(10)	175.1(10)
	172.5(8)	173.7(11)	169.8(10)	176.4(10)

- a) Estimated standard deviations in the least significant figures are given in parentheses. Phenyl groups were refined as rigid groups with C-C as 1.39Å and C-H as 1.00Å.
- b) See caption to Figure 3. These distances are for CN-CN linking monomers 1-1, 2-2, etc.
- c) These distances are for CN-CN linking monomers 1 and 2, 3 and 4.
- d) Because the phenyl rings were refined as rigid groups errors in distances involving them are not available.

TABLE III

Crystal Data for [closo-3-(PPh<sub>3</sub>)-3,3-[C(Ph)-C(PPh<sub>3</sub>)-C(H)-C(Ph)]-3,1,2-Rh-C<sub>2</sub>B<sub>9</sub>H<sub>11</sub>](5) and [closo-3-(PPh<sub>3</sub>)-3-(μ-CN)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>]<sub>4</sub>·5C<sub>6</sub>H<sub>6</sub> (7).

COMPOUND		(5)	(7)
Crystal System		Triclinic, $P\bar{1}$ , Z=2	Monoclinic, P2/a, Z=4
Cell	a	12.763(6) Å	26.046 (8) Å
	b	13.348(5) Å	15.626 (3) Å
	c	14.561(7) Å	30.355 (8) Å
	α	91.58 (3)°	90.
	β	93.72 (3)°	106.71 (2)°
	γ	74.64 (3)°	90.
	v	2404 Å <sup>3</sup>	11833 Å <sup>3</sup>
Calculated density		1.34g/cm <sup>3</sup> (119°K)	1.35g/cm <sup>3</sup> (119°K)
Observed density		-----	1.24g/cm <sup>3</sup>
Sinθ/λ data collection limit		0.538 Å <sup>-1</sup>	0.538 Å <sup>-1</sup>
Scan rate		2°/min.	3°/min.
Scan range		(2.2 + 0.692 tan θ)°	(1.8 + 0.688 tan θ)°
Background		20s	0.5 x scan time
Absorption coefficient		4.56 cm <sup>-1</sup> (MoKα)	5.77 cm <sup>-1</sup> (MoKα)
Number of reflections collected		6266 (hemisphere)	15992 (quadrant)
Number of reflections used in analysis		4493	9732
Agreement factors	R	0.048	0.063
	Rw	0.051	0.078
Goodness-of-fit parameter		1.50	2.03
Data-to-parameter ratio		26	20.

TABLE IV

A) Positional (in fractional coordinates) and Thermal Parameters<sup>a</sup> of the Non-GroupAtoms in [closo-3-(PPh<sub>3</sub>)-3,3-{C(Ph)-C(PPh<sub>3</sub>)-C(H)-C(Ph)}-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>](5).

Atom	x	y	z	$\frac{\sigma^2}{B(A)}$
Rh	.15780(4)	.15338(4)	.28636(3)	b
P(1)	.0143(1)	.3009(1)	.2556(1)	b
P(2)	.4413(1)	.2716(1)	.1911(1)	b
C(1)	.0982(5)	.0544(4)	.3857(4)	1.5(1)
C(2)	.0479(5)	.0430(5)	.2832(4)	1.8(1)
B(4)	.2362(5)	.0269(5)	.3874(4)	1.6(1)
B(5)	.1683(6)	-.0622(5)	.4300(5)	2.1(1)
B(6)	.0449(6)	-.0506(6)	.3638(5)	2.0(1)
B(7)	.1426(6)	.0042(5)	.2060(5)	1.6(1)
B(8)	.2712(6)	-.0083(5)	.2718(5)	1.4(1)
B(9)	.2772(6)	-.1065(6)	.3576(5)	2.0(1)
B(10)	.1592(6)	-.1539(6)	.3421(5)	2.0(1)
B(11)	.0762(6)	-.0856(6)	.2496(5)	1.9(1)
B(12)	.2194(6)	-.1189(6)	.2444(5)	1.9(1)
C(3)	.2348(5)	.2403(4)	.3721(4)	1.2(1)
C(4)	.3069(5)	.2789(5)	.3302(4)	1.4(1)
C(5)	.3173(4)	.2559(4)	.2325(4)	1.3(1)
C(6)	.2492(5)	.2016(4)	.1929(4)	1.3(1)
H(1)	.056(3)	.101(3)	.429(3)	1.6(9)
H(2)	-.031(3)	.084(3)	.266(3)	0.6(8)
H(4)	.280(3)	.053(3)	.438(3)	1.6(8)
H(5)	.168(4)	-.071(3)	.507(3)	1.6(10)
H(6)	-.025(4)	-.052(4)	.393(3)	1.3(10)
H(7)	.125(4)	.020(4)	.137(4)	2.1(12)
H(8)	.344(4)	-.007(4)	.245(3)	2.0(11)
H(9)	.357(4)	-.149(4)	.383(3)	1.3(10)
H(10)	.156(4)	-.235(5)	.358(3)	1.2(10)
H(11)	.014(5)	-.116(5)	.204(4)	3.3(15)
H(12)	.254(5)	-.176(5)	.196(4)	3.2(15)
HC4	.356(4)	.314(4)	.366(3)	1.5(11)

Table IV (continued)

B) Anisotropic Thermal Parameters<sup>c</sup>

Atom	$10^5 \beta_{11}$	$10^5 \beta_{22}$	$10^5 \beta_{33}$	$10^5 \beta_{12}$	$10^5 \beta_{13}$	$10^5 \beta_{23}$
Rh	217(3)	126(3)	118(2)	-40(2)	9(2)	-10(2)
P(1)	226(11)	169(10)	144(8)	-56(8)	-2(7)	-20(8)
P(2)	213(11)	232(11)	198(8)	-57(9)	20(8)	13(7)

C) Rigid Group Parameters<sup>d</sup>

Group	$\underline{x}$	$\underline{y}$	$\underline{z}$	$\underline{\Phi}$	$\underline{\Theta}$	$\underline{\rho}$	$B(\text{\AA}^2)$
Phen 1	.2259(3)	.2510(3)	.4750(2)	-1.684(2)	-3.046(2)	1.673(2)	2.11(5)
Phen 2	.2424(3)	.1828(3)	.0900(2)	0.020(2)	3.021(2)	-1.607(2)	2.05(5)
Phen 3	-.0867(3)	.3419(3)	.3466(2)	-.421(2)	-3.050(2)	2.363(2)	1.76(5)
Phen 4	-.0810(3)	.2896(3)	.1569(2)	-2.526(2)	2.765(2)	-1.016(2)	1.99(5)
Phen 5	.0483(3)	.4234(2)	.2296(2)	1.590(5)	1.912(2)	-.508(5)	1.64(5)
Phen 6	.4565(3)	.3960(2)	.2325(3)	-2.006(2)	-2.795(2)	2.744(2)	2.12(5)
Phen 7	.5503(3)	.1676(3)	.2414(3)	-1.251(3)	2.552(2)	0.511(3)	2.59(5)
Phen 8	.4594(3)	.2660(3)	.0634(2)	3.092(2)	-3.103(2)	-1.778(2)	2.09(5)

a) Estimated standard deviations in the least significant figures are given in parentheses.

b) Anisotropic thermal parameters.

c) The form of the anisotropic thermal ellipsoid is:

$$\exp[(\beta_{11}h^2 + \beta_{22}k^2 + \beta_{33}l^2 + 2\beta_{12}hk + 2\beta_{13}hl + 2\beta_{23}kl)]$$

d)  $x$ ,  $y$  and  $z$  are the fractional coordinates of the origin of the rigid group and  $\phi$ ,  $\theta$  and  $\rho$  the rotation angles in radians, as defined by C. Scheringer, Acta Crystallogr., 16, 546 (1963).

Table V

Positional<sup>a</sup> (in fractional coordinates) and Thermal Parameters for  
 $[\{\text{closo-3-(PPh}_3\text{)-3-(}\mu\text{-CN)-3,1,2-RhC}_2\text{B}_9\text{H}_{11}\}_4 \cdot 5\text{C}_6\text{H}_6](7)$ .

Nongroup Atoms	$10^4x$	$10^4y$	$10^4z$	$\frac{\sigma^2}{B(\text{\AA})}$
Rh(1)	1741.3(3)	-713.3(5)	4218.3(3)	b
P(1)	2355(1)	-1109(2)	3839(1)	b
CN(1)	1939(4)	537(6)	4184(3)	1.4(2)
CN(1)'	2342(4)	-847(5)	4820(3)	1.4(2)
C(11)	1167(4)	-1063(7)	4606(4)	2.3(3)
C(12)	964(5)	-217(7)	4273(4)	2.5(3)
B(14)	1322(5)	-1902(7)	4298(4)	1.2(2)
B(15)	719(5)	-1839(8)	4456(4)	1.7(2)
B(16)	500(5)	-813(8)	4444(4)	1.8(2)
B(17)	968(6)	-471(8)	3695(5)	1.7(3)
B(18)	1189(6)	-1568(8)	3712(4)	1.5(3)
B(19)	714(6)	-2206(8)	3895(4)	1.9(3)
B(110)	209(5)	-1538(8)	3979(4)	2.0(2)
B(111)	362(6)	-452(9)	3861(5)	1.7(3)
B(112)	499(6)	-1316(8)	3526(4)	1.9(3)
Rh(2)	2251.0(3)	2490.3(5)	4098.6(3)	b
P(2)	1366(1)	2934(2)	3948(1)	b
CN(2)	2006(4)	1259(5)	4153(3)	1.3(2)
CN(2)'	2429(4)	2558(6)	4802(3)	1.7(2)
C(21)	2796(4)	1943(7)	3709(4)	1.9(2)
C(22)	2271(5)	2491(8)	3378(4)	1.8(2)
B(24)	3129(5)	2501(8)	4166(4)	2.0(3)
B(25)	3401(6)	2333(8)	3709(4)	2.5(3)
B(26)	2861(5)	2320(8)	3203(4)	2.1(2)
B(27)	2273(5)	3542(7)	3622(4)	1.6(2)
B(28)	2827(5)	3539(7)	4152(4)	1.8(2)
B(29)	3428(6)	3358(9)	3974(4)	2.1(3)
B(210)	3245(6)	3262(9)	3359(5)	2.3(3)
B(211)	2545(6)	3350(9)	3147(5)	2.3(3)
B(212)	2899(5)	4004(8)	3624(4)	1.9(3)

Table V (continued)

Rh(3)	3050.1(3)	5805.2(5)	867.8(3)	b
P(3)	2366(1)	6259(2)	1154(1)	b
CN(3)	2604(4)	5898(6)	189(3)	1.7(2)
CN(3)'	2748(4)	4594(6)	860(3)	1.6(2)
C(31)	3864(4)	5207(7)	1134(4)	2.5(2)
C(32)	3792(5)	5724(7)	655(4)	2.2(2)
B(34)	3690(5)	5802(8)	1536(4)	2.0(3)
B(35)	4370(6)	5579(10)	1589(5)	2.6(3)
B(36)	4435(6)	5517(9)	1022(5)	3.1(3)
B(37)	3573(6)	6800(9)	708(5)	2.3(3)
B(38)	3518(5)	6867(8)	1298(4)	1.4(3)
B(39)	4172(5)	6631(8)	1676(4)	2.4(2)
B(310)	4634(6)	6470(9)	1350(5)	2.7(3)
B(311)	4260(7)	6540(10)	755(4)	2.9(3)
B(312)	4104(6)	7253(8)	1165(4)	2.3(3)
Rh(4)	2475.6(3)	2609.8(5)	859.2(3)	b
P(4)	3369(1)	2250(2)	1074(1)	b
CN(4)	2649(4)	3885(6)	844(3)	1.7(2)
CN(4)'	2500(4)	2504(5)	189(3)	2.0(2)
C(41)	1695(5)	2971(7)	982(4)	2.2(2)
C(42)	1623(5)	2152(7)	629(4)	2.2(2)
B(44)	2216(5)	2811(8)	1479(4)	1.5(2)
B(45)	1531(6)	2729(8)	1467(4)	2.2(3)
B(46)	1147(6)	2317(9)	931(5)	2.4(3)
B(47)	2105(5)	1339(8)	885(4)	1.7(2)
B(48)	2472(6)	1765(8)	1445(4)	1.9(3)
B(49)	2006(6)	1947(9)	1759(5)	2.2(3)
B(410)	1356(6)	1611(10)	1418(5)	2.6(3)
B(411)	1418(5)	1260(8)	878(4)	2.2(3)
B(412)	1928(6)	1043(9)	1387(5)	2.1(3)

### Anisotropic Thermal Parameters

<u>Atom</u>	<u><math>10^5 \beta_{11}</math></u>	<u><math>10^5 \beta_{22}</math></u>	<u><math>10^5 \beta_{33}</math></u>	<u><math>10^5 \beta_{12}</math></u>	<u><math>10^5 \beta_{13}</math></u>	<u><math>10^5 \beta_{23}</math></u>
Rh(1)	35(2)	102(4)	36(1)	-4(2)	5(1)	2(2)
P(1)	63(6)	146(12)	61(4)	8(7)	28(4)	10(6)
Rh(2)	50(2)	107(4)	32(1)	-3(2)	11(1)	6(2)
P(2)	57(6)	117(12)	50(4)	7(7)	9(4)	14(5)
Rh(3)	62(2)	119(4)	46(1)	-11(2)	18(1)	-6(2)
P(3)	81(6)	128(12)	59(4)	10(7)	29(4)	1(6)
Rh(4)	71(2)	110(4)	42(1)	-6(2)	16(1)	3(2)
P(4)	69(6)	155(13)	55(4)	5(7)	16(4)	-5(5)

### Rigid Group Parameters<sup>d</sup>

<u>Name</u>	<u><math>10^4 x</math></u>	<u><math>10^4 y</math></u>	<u><math>10^4 z</math></u>	<u>Phi</u>	<u>Theta</u>	<u>Rho</u>	<u><math>B(A^2)</math></u>
Phenyl 1	2596(3)	-2210(3)	3984(3)	2.067(4)	2.533(3)	2.762(5)	2.5(1)
Phenyl 2	2108(3)	-974(3)	3219(2)	.160(3)	-2.939(2)	-1.594(5)	2.4(1)
Phenyl 3	2979(3)	-499(4)	3971(3)	-2.403(7)	2.182(3)	2.763(7)	3.0(1)
Phenyl 4	1718(2)	5700(4)	960(2)	2.900(6)	-2.102(3)	-.682(7)	2.3(1)
Phenyl 5	2141(3)	7344(3)	1020(3)	1.720(4)	-2.563(2)	-.313(5)	2.4(1)
Phenyl 6	2578(3)	6149(3)	1779(2)	-3.118(3)	3.039(2)	1.611(5)	2.3(1)
Phenyl 7	3731(3)	2507(4)	1667(2)	.595(3)	3.026(3)	1.234(5)	2.6(1)
Phenyl 8	3444(3)	1121(3)	963(3)	2.418(9)	-1.954(3)	-2.373(9)	3.0(1)
Phenyl 9	3798(3)	2760(4)	766(3)	-2.329(5)	-2.557(3)	-2.518(6)	3.4(1)
Phenyl 10	925(3)	2402(5)	4234(2)	2.332(7)	2.109(3)	1.245(9)	2.8(1)
Phenyl 11	1270(3)	4028(3)	4107(2)	1.656(4)	2.631(2)	.362(4)	2.1(1)
Phenyl 12	1034(3)	2796(3)	3337(2)	-3.096(3)	-2.966(2)	-1.460(5)	2.4(1)

a) Standard deviations are given in parentheses.

b) Anisotropic thermal parameters.

c) The form of the anisotropic thermal ellipsoid is:

$$\exp[-(\beta_{11}h^2 + \beta_{22}k^2 + \beta_{33}l^2 + 2\beta_{12}hk + 2\beta_{13}hl + 2\beta_{23}kl)]$$

d) x, y, and z are the fractional coordinates of the group origin, and phi, theta, and rho the rotation angles in radians, as defined by

C. Scheringer, Acta Crystallogr., **16**, 546 (1963).

Figure 1.

Structure of [closo-3-(PPh<sub>3</sub>)-3,3-{C(Ph)-C(PPh<sub>3</sub>)-C(H)-C(Ph)}-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>](5).

The phenyl rings and carborane hydrogens have been omitted for clarity. The thermal ellipsoids represent 50% probability.






Figure 2. Proposed bonding for the  
metallocycle (5).







Figure 3.

One of the two crystallographically unique molecules of [closo-3-(PPh<sub>3</sub>)-3-(μ-CN)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>]<sub>4</sub>(7), with the phenyl rings, and hydrogen atoms omitted for clarity and the atoms of the C<sub>2</sub>B<sub>9</sub> cages drawn with an artificially small radius. The asymmetric unit of the crystal unit cell consists of two distinct one-half tetramers. The other half of each tetramer is generated by a twofold rotation axis passing through the center of two opposing C≡N bonds. The consequent 50%-50% disorder in these CN linkages indicates the other CN ligands suffer from 50%-50% disorder also, assuming that each Rh is bound to one N and one C atom of two bridging CN<sup>-</sup> ligands.






SCHEME I




SCHEME I







SCHEME III



(i) Hydrochloric acid, THF

(ii) Triphenylphosphine, THF

(iii) Hydrochloric acid, Et<sub>2</sub>O.

(iv) Triphenylphosphine, Et<sub>2</sub>O.

Reactions (i) through (iv) proceed at room temperature.

## SCHEME IV

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER TECHNICAL REPORT 116	2. GOVT ACCESSION NO. AD A113 112	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) "Reactions at the Metal Vertex of a Monometal Metallocarborane Cluster. The Chemistry of [closo-3,3-(PPh <sub>3</sub> ) <sub>2</sub> -3-(HSO <sub>3</sub> ) <sub>2</sub> -3,1,2-RhC <sub>2</sub> B <sub>9</sub> H <sub>11</sub> ] And [closo-3-(PPh <sub>3</sub> ) <sub>2</sub> -3,3-(NO <sub>3</sub> ) <sub>2</sub> -3,1,2-RhC <sub>2</sub> B <sub>9</sub> H <sub>11</sub> ]."		5. TYPE OF REPORT & PERIOD COVERED Interim
7. AUTHOR(s) W.C. Kalb, Z. Demidowicz, D.M. Speckman, Carolyn Knobler, Raymond G. Teller and M. Frederick Hawthorne		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS The University of California Department of Chemistry Los Angeles, California 90024		8. CONTRACT OR GRANT NUMBER(s) N00014-76-C-0390
11. CONTROLLING OFFICE NAME AND ADDRESS Chemistry Branch Office of Naval Research Washington, D.C. 20360		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 053-608
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE March 9, 1982
		13. NUMBER OF PAGES
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) <div style="border: 1px solid black; padding: 5px; text-align: center;">This document has been approved for public release and sale; its distribution is unlimited.</div>		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)		

# ABSTRACT

Reaction of [closo-3,3-(PPh<sub>3</sub>)<sub>2</sub>-3-(H)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>](1) with sulfuric or nitric acid affords [closo-3,3-(PPh<sub>3</sub>)<sub>2</sub>-3-(HSO<sub>4</sub>)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>](2) or [closo-3-(PPh<sub>3</sub>)-3,3-(NO<sub>3</sub>)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>](3), respectively. Compound (3) can also be prepared from nitric acid and the dimeric metallocarborane [closo-Rh(PPh<sub>3</sub>)(C<sub>2</sub>B<sub>9</sub>H<sub>11</sub>)]<sub>2</sub> or from NO<sub>2</sub>/N<sub>2</sub>O<sub>4</sub> and (1). Complexes (2) and (3) have been used to prepare other new metallocarboranes; namely, [closo-3-(PPh<sub>3</sub>)-3,3-{C(Ph)-C(PPh<sub>3</sub>)-C(H)-C(Ph)}-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>](5), [{closo-3-(PPh<sub>3</sub>)-3-(μ-CN)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>}]<sub>4</sub>(7), [closo-3-(PPh<sub>3</sub>)-3-(L)-3-(NO<sub>3</sub>)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub> (L = CO, (8); L = PPh<sub>3</sub>, (9)), [closo-3,3-(PMe<sub>2</sub>Ph)<sub>2</sub>-3-NO<sub>3</sub>-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>](10) and [closo-3-(PPh<sub>3</sub>)-3-(CO)-3-(Cl)-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>](11). Complexes (5) and (7) have been characterized by X-ray crystallography. The reactions of these new metallocarboranes described herein are representative of interconversions carried out at a discrete transition metal vertex of a cluster species.

Complex (5) crystallizes in space group P $\bar{1}$  with 2 formula units in a cell of dimensions  $a = 12.763(6)$  Å,  $b = 13.348(5)$  Å,  $c = 14.561(7)$  Å,  $\alpha = 91.58(3)^\circ$ ,  $\beta = 93.72(3)^\circ$ , and  $\gamma = 74.64(3)^\circ$ . Data were collected at  $-154^\circ$  C on a Picker FACS-1 diffractometer using the  $\theta$ - $2\theta$  scan method. Least-squares refinement, including anisotropic vibration parameters for Rh and P, isotropic vibration parameters for other nonhydrogen atoms, and with each phenyl group described as a rigid group having a single isotropic vibration parameter, led to final conventional agreement indices (on F) of  $R = 0.048$  and  $R_w = 0.051$ , based on 4493 unique reflections having  $I > 3\sigma(I)$ . The molecule consists of a [C<sub>2</sub>B<sub>9</sub>H<sub>11</sub>]<sup>2-</sup> cage and a triphenylphosphine ligand bound to the metal atom of the trisubstituted metallopentacycle Rh-C(Ph)-C(PPh<sub>3</sub>)-C(H)-C(Ph). Rh-C, Rh-B, B-B, B-C, and C-C distances are normal for a 3,1,2-RhC<sub>2</sub>B<sub>9</sub> closo-rhodacarborane fragment and the pattern of short-long-short C-C bond lengths in the RhC<sub>4</sub> ring is reminiscent of

-1a-

a penta-substituted cis-butadiene.

Complex (7)  $\cdot 5C_6H_6$  crystallizes in space group  $P2_1/a$  with 4 formula units in a cell of dimensions  $a = 26.046(8) \text{ \AA}$ ,  $b = 15.626(3) \text{ \AA}$ ,  $c = 30.355(8) \text{ \AA}$ , and  $\beta = 106.71(2)^\circ$ . Data were collected at  $-154^\circ \text{ C}$  on a Syntex P1 diffractometer using the  $\Theta$ - $2\Theta$  scan method. Least-squares refinement, including vibration parameters and rigid group assignments as described above, led to final conventional agreement indices (on F) of  $R = 0.063$  and  $R_w = 0.078$ , based on 9732 unique reflections have  $I > 3\sigma(I)$ . The molecules consist of four discrete closo-phosphino-rhodacarborane moieties joined together through their respective metal vertices by cyano ligand bridges. Each tetramer possesses a crystallographic 2-fold axis; the 2 non-crystallographically equivalent tetramers are very similar. The Rh-Rh separation is approximately  $5 \text{ \AA}$ . Bond distances within each icosahedral fragment are normal for such a closo-Rh(III)-metallo-carborane.

APPENDIX

TECHNICAL REPORT DISTRIBUTION LIST, 053

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. R.N. Grimes Department of Chemistry University of Virginia Charlottesville, Virginia 22901	1	Dr. M.H. Chisholm Department of Chemistry Indiana University Bloomington, Indiana 47401	1
Dr. D.B. Brown Department of Chemistry University of Vermont Burlington, Vermont 05401	1	Dr. B. Foxman Department of Chemistry Brandeis University Waltham, Massachusetts 02154	1
Dr. W.B. Fox Chemistry Division Naval Research Laboratory Code 6130 Washington, D.C. 20375	1	Dr. T. Marks Department of Chemistry Northwestern University Evanston, Illinois 60201	1
Dr. J. Adcock Department of Chemistry University of Tennessee Knoxville, Tennessee 37916	1	Dr. G. Geoffrey Department of Chemistry Pennsylvania State University University Park, Pennsylvania 16802	1
Dr. A. Cowley Department of Chemistry University of Texas Austin, Texas 78712	1	Dr. J. Zuckerman Department of Chemistry University of Oklahoma Norman, Oklahoma 73019	1
Dr. W. Hatfield Department of Chemistry University of North Carolina Chapel Hill, North Carolina 27514	1	Professor O.T. Beachley Department of Chemistry State University of New York Buffalo, New York 14214	1
Dr. D. Seyferth Department of Chemistry Massachusetts Institute of Technology Cambridge, Massachusetts 02139	1	Professor P.S. Skell Department of Chemistry The Pennsylvania State University University Park, Pennsylvania 16802	1
		Professor K.M. Nicholas Department of Chemistry Boston College Chestnut Hill, Massachusetts 02167	1
Professor H. Abrahamson Department of Chemistry University of Oklahoma Norman, Oklahoma 73019	1	Professor R. Neilson Department of Chemistry Texas Christian University Fort Worth, Texas 76129	1
Professor M. Newcomb Texas A&M University Department of Chemistry College Station, Texas 77843	1	Professor Richard Eisenberg Department of Chemistry University of Rochester Rochester, New York 14627	1

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
Office of Naval Research Attn: Code 472 800 North Quincy Street Arlington, Virginia 22217	2	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 1211 Research Triangle Park, N.C. 27709	1
ONR Branch Office Attn: Dr. George Sandoz 536 S. Clark Street Chicago, Illinois 60605	1	Naval Ocean Systems Center Attn: Mr. Joe McCartney San Diego, California 92152	1
	1	Naval Weapons Center Attn: Dr. A.B. Amster, Chemistry Division China Lake, California 93555	1
ONR Branch Office 1030 East Green Street Pasadena, California 91106	1	Naval Civil Engineering Laboratory Attn: Dr. R.W. Drisko Port Hueneme, California 93401	1
ONR Branch Office Attn: Dr. L.H. Peebles Building 114, Section D 666 Summer Street Boston, Massachusetts 02210	1	Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
Director, Naval Research Laboratory Attn: Code 6100 Washington, D.C. 20390	1	Dr. A.L. Slafkosky Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
The Assistant Secretary of the Navy (R,E&S) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1	Office of Naval Research Attn: Dr. Richard S. Miller 800 N. Quincy Street Arlington, Virginia 22217	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Department of the Navy Washington, D.C. 20360	1	Naval Ship Research and Development Center Attn: Dr. G. Bosmajian, Applied Chemistry Division Annapolis, Maryland 21401	1
Defense Documentation Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Naval Ocean Systems Center Attn: Dr. S. Yamamoto, Marine Sciences Division San Diego, California 91232	1
Dr. Fred Saalfeld Chemistry Division Naval Research Laboratory Washington, D.C. 20375	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1



P4-2/A25

472:GAN:716:tam  
78u472-608

TECHNICAL REPORT DISTRIBUTION LIST, GEN

No.  
Copies

Dr. Rudolph J. Marcus  
Office of Naval Research  
Scientific Liaison Group  
American Embassy  
APO San Francisco 96503

1

Mr. James Kelley  
DTNSRDC Code 2803  
Annapolis, Maryland 21402

1

P4-2/A25

472:GAN:J16:tam  
78u472-608

TECHNICAL REPORT DISTRIBUTION LIST, GEN

No.  
Copies

Dr. Rudolph J. Marcus  
Office of Naval Research  
Scientific Liaison Group  
American Embassy  
APO San Francisco 96503

1

Mr. James Kelley  
DTNSRDC Code 2803  
Annapolis, Maryland 21402

1

DATE  
FILMED  
—8